

Research Group: Public Economics

May, 2012

"Euclidean Preferences, Option Sets and Strategy Proofness"

*Georges Bordes
Gilbert Laffond
Michel Le Breton*

Euclidean Preferences, Option Sets and Strategy Proofness*

Georges Bordes[†] Gilbert Laffond[‡] Michel Le Breton[§]

May 2012[¶]

Abstract

In this note, we use the technique of option sets to sort out the implications of coalitional strategyproofness in the spatial setting. We also discuss related issues and open problems.

1 Introduction

The purpose of this note is to illustrate the power of an ingenious technique pioneered independently by Barbera and Peleg (1990) and Laffond (1980) and called the *technique of option sets*¹ which has been used extensively and successfully to characterize the implications of strategyproofness in many different settings. To quote Zhou (1991), one, out of many scholars, who has used it: "It is direct and simple, invoking neither the Arrow theorem, nor any monotonicity argument. Yet, it is so powerful that under its framework many interesting issues can be adressed"². Out of many contributions², we can cite for instance Barbera, Masso and Neme (1997) and Barbera, Masso and Serizawa (1998) who use the technique of option

*We are grateful to Salvador Barbera for the many stimulating scientific conversations we had over years and very pleased to offer him this little contribution which has remained unpublished. The last author would like to point out that his own interest for strategyproofness has been very much influenced by his very nice 1990 paper with Bezalel Peleg.

[†]Our dear friend Georges Bordes passed away in 2005. He was a very good person, a talented scholar and a friend of Salvador. He would have been very happy to offer this little present to his friend.

[‡]Laboratoire d'Econométrie, CNAM, Paris.

[§]Toulouse School of Economics and Institut Universitaire de France.

[¶]We would like to thank an anonymous referee for an extremely careful reading of an earlier version of this manuscript. We would also like to thank Jérôme Renault and John Weymark for a very useful correspondence.

¹Border and Jordan (1983) call them manipulation sets. These sets also appear in Laffont (1987).

²I cannot cite here all the contributions who use that technique as they are too many.

sets to investigate the case in which the domain is the class of multidimensional single-peaked preferences and the range is a compact set and Berga and Serizawa (2000) who use it in their exploration of maximal domains for strategy-proofness.

This simple notion is defined as follows. Consider a society of individuals who has to select an alternative out of a feasible set X and a social choice mechanism mapping any conceivable profile R of preferences into an alternative. The option set $O_S(R_{-S})$ of a coalition S of individuals (given the preferences R_{-S} reported by the individuals outside the coalition) is the set of alternatives that they can reach through an appropriate joint report R_S of their preferences. Therefore, the option set of coalition S describes the "scope of influence" of coalition S given the profile R_{-S} of reports by individuals outside S . A social environment is characterized by a set X and a domain \mathcal{D} of admissible preferences. Given a social environment (X, \mathcal{D}) , the technique of options sets consists in sorting out gradually the properties that these different option sets must possess if the mechanism is required to satisfy some properties, on top of which some weak or strong versions of strategyproofness.

In this note, we illustrate this technique in the case where $X = \mathbb{R}^m$ and \mathcal{D} is the set of Euclidean preferences. This result is extracted from a work that was done twenty years ago³ (Bordes, Laffond and Le Breton (1990)). The result states that, if $m \geq 2$, any surjective and coalitional strategyproof mechanism is dictatorial. The main part of the proof consists in the analysis of the case where there are two dimensions and two individuals. It uses elementary tools from basic geometry and is self contained. Most of the proof consists in showing that to prevent the mechanism to be manipulated by any one of the two individuals, say 2, the option set of 2 must be a disk. This proof is twenty years old and, likely, some better self-contained proofs can be provided. Since then, many new results have been discovered. In the last section, we offer a brief account of some of the main achievements and open problems in this branch of the literature.

2 The Model

We consider a society $N = \{1, 2, \dots, n\}$ of individuals who has to select an alternative in the Euclidean space $X = \mathbb{R}^m$. The preference R_i of any individual $i \in N$ is entirely described by a point $p_i \in \mathbb{R}^m$ and defined as follows : xR_iy iff $\|x - p_i\| \leq \|y - p_i\|$ where $\|\cdot\|$ denotes the usual Euclidean norm on \mathbb{R}^m ⁴. The point p_i is called the peak of individual i as it is

³The old version was containing a brief account of some of the results contained in the doctoral thesis of the second author. In the last part of this note, we offer a short presentation of some of his ideas.

⁴ $\|x\| = \langle x, x \rangle^{\frac{1}{2}}$ where $\langle \cdot, \cdot \rangle$ denotes the usual inner product.

the (unique) mostly preferred alternative of individual i : any individual orders alternatives according to their distance with respect to his peak. If an alternative is interpreted as the location of a public facility and p_i as the place of residence of individual i , Euclidean preferences can be interpreted as preferences for smallest distances to the facility when the transportation network is unrestricted. Hereafter, we will denote by \mathcal{E} the domain of Euclidean preferences over \mathbb{R}^m .

A social choice mechanism is a mapping C from \mathcal{E}^n into X . Since preferences in \mathcal{E} are identified to their peaks, we will alternatively describe C as a mapping from \mathbb{R}^{nm} into \mathbb{R}^m mapping a profile $\mathbf{p} = (p_1, p_2, \dots, p_n)$ of peaks into a social alternative $C(\mathbf{p}) = (C^1(\mathbf{p}), C^2(\mathbf{p}), \dots, C^m(\mathbf{p}))$. Given a profile $\mathbf{p} \in \mathbb{R}^{nm}$ and a coalition $S \subseteq N$, we denote by $\mathbf{p}_{-S} \in (\mathbb{R}^m)^{N \setminus S}$ the restriction of the profile to $N \setminus S$. A social choice mechanism C is strategy-proof if there does not exist $i \in N$, $\mathbf{p} \in \mathbb{R}^{nm}$ and $\hat{p}_i \in \mathbb{R}^m$ such that $\|C(p_{-i}, \hat{p}_i) - p_i\| < \|C(\mathbf{p}) - p_i\|$. A mechanism is coalitional strategy-proof if there does not exist $S \subseteq N$, $\mathbf{p} \in \mathbb{R}^{nm}$ and $\hat{\mathbf{p}}_S \in (\mathbb{R}^m)^S$ such that $\|C(\mathbf{p}_{-S}, \hat{\mathbf{p}}_S) - p_i\| < \|C(\mathbf{p}) - p_i\|$ for all $i \in S$. The range of the mechanism C is the set $R(C) \equiv \{x \in \mathbb{R}^m : x = C(\mathbf{p}) \text{ for some } \mathbf{p} \in \mathbb{R}^{nm}\}$. The mechanism C is unanimous if $C(p, p, \dots, p) = p$. C is anonymous if $C(p_1, p_2, \dots, p_n) = C(p_{\sigma(1)}, p_{\sigma(2)}, \dots, p_{\sigma(n)})$ for all permutations $\sigma : N \rightarrow N$ and all $\mathbf{p} \in \mathbb{R}^{nm}$. C is (component-wise) continuous if each component C^j is continuous. Finally, a social choice mechanism C is dictatorial if there exists $i \in N$ such that $C(\mathbf{p}) = p_i$ for all $\mathbf{p} \in \mathbb{R}^{nm}$.

3 The Result

Proposition If $m \geq 2$, then any coalitional strategy-proof social choice mechanism C over the domain \mathcal{E} such that $R(C) = \mathbb{R}^m$ is dictatorial .

4 The Proof

As already pointed out, the proof is based on the technique of options sets. Given a profile $\mathbf{p} \in \mathbb{R}^{nm}$ and a coalition $S \subseteq N$, we define the option set of coalition S at profile \mathbf{p} denoted $O_S(\mathbf{p})$ as the set $\left\{x \in \mathbb{R}^m : x = C(\mathbf{p}_{-S}, \hat{\mathbf{p}}_S) \text{ for some } \hat{\mathbf{p}}_S \in (\mathbb{R}^m)^S\right\}$. It will be often denoted $O_S(\mathbf{p}_{-S})$ as it only depends upon the subprofile \mathbf{p}_{-S} . When $S = \{i\}$ for some $i \in N$, $O_S(\mathbf{p})$ will be denoted $O_i(\mathbf{p}_{-i})$. The option set of a coalition S , given the preferences reported by the individuals outside the coalition, is the set of alternatives attainable by the members of S if they jointly control the preferences that they report. It is the range of the social choice mechanism $C_S(\mathbf{p}_{-S})$ defined on $(\mathbb{R}^m)^S$. We note that $O_N(\mathbf{p}) = R(C)$.

The following two properties are quite general (i.e. independent of the specific Euclidean setting considered here) and have been demonstrated several times. We will not repeat these arguments here.

Step 1 : If $p \in R(C)$, then $C(p, p, \dots, p) = p$

Step 2 (Closeness) : For all $\mathbf{p} \in \mathbb{R}^{nm}$ and $S \subseteq N$, $O_S(\mathbf{p}_{-S})$ is a closed subset of \mathbb{R}^m

4.1 The Case where $n=m=2$

In this section we will denote simply $O_1(p_2)$ and $O_2(p_1)$ the options set of individuals 1 and 2. Step after step, we will refine our knowledge of the options sets.

Step 3 : $p_1 \in O_2(p_1)$ and $p_2 \in O_1(p_2)$

Step 4 : $\|C(p_1, p_2) - p_2\| \leq \|p'_2 - p_2\|$ for all $p'_2 \in O_2(p_1)$ and $\|C(p_1, p_2) - p_1\| \leq \|p'_1 - p_1\|$ for all $p'_1 \in O_1(p_2)$.

Step 5 (Efficiency) : $C(p_1, p_2) \in [p_1, p_2]$

Step 6 (Star-shapedness) : $O_2(p_1)$ is star-shaped with respect to p_1 .

If $O_2(p_1) = \{p_1\}$ the conclusion follows. Suppose that there exists $t \neq p_1$ such that $t \in O_2(p_1)$. We want to show that $[p_1, t] \subseteq O_2(p_1)$. Let $z \in [p_1, t]$ and assume on the contrary that $z \notin O_2(p_1)$. Since $O_2(p_1)$ is closed, so is $E \equiv [z, t] \cap O_2(p_1)$. This implies that the program $\underset{w \in E}{\text{Min}} \|w - z\|$ has a solution, say v . By construction: $[z, v] \cap O_2(p_1) = \emptyset$. Let $p_2 \in]\frac{z+v}{2}, v[$. Since $]\frac{z+v}{2}, v[\cap O_2(p_1) = \emptyset$, we deduce from step 5 that $C(p_1, p_2) \in [p_1, z[$. Since $\|v - p_2\| < \|z - p_2\|$, this contradicts step 4.

An immediate consequence of step 6 is the following. Given any ray L with endpoint p_1 , if $L \cap O_2(p_1)$ is bounded, then there exists $z = z(L)$ such that $L \cap O_2(p_1) = [p_1, z]$.

Step 7 (Tangency) : Let L be a ray with endpoint p_1 and $L^\perp(z)$ be the line orthogonal to L containing z . Then $O_2(p_1)$ is contained in the half plane $P(z)$ with frontier $L^\perp(z)$ and containing p_1 i.e. $\langle x - z, p_1 - z \rangle \geq 0$ for all $x \in O_2(p_1)$.

Assume on the contrary that there exists $x \in O_2(p_1)$ such that:

$$\langle x - z, p_1 - z \rangle < 0$$

Then, there exists $y \in L$ such that $\|y - x\| < \|y - z\|$. The existence of such y follows from the fact that if we take y on L sufficiently far from z , then x will be in the interior of the disk centered on y with radius $\|y - z\|$. This argument is illustrated on figure 1.

Insert Figure 1 here

From steps 4 and 5, $C(p_1, y) = z$ but since $x \in O_2(p_1)$, there exists p_2 such that $x = C(p_1, p_2)$. Since $\|y - C(p_1, p_2)\| < \|y - C(p_1, y)\|$, this contradicts strategyproofness.

Step 8 : If $O_2(p_1) \cap L$ is bounded for some L , then $O_2(p_1)$ is bounded.

The claim follows from a repeated application of step 7.

Step 9 : If $O_2(p_1) \cap L = \{p_1\}$ for some L , then $O_2(p_1) = \{p_1\}$.

As for step 8, it follows from a repeated application of step 7.

At this stage, for every $p_1 \in \mathbb{R}^2$, there are three possible cases:

Case 1 : $O_2(p_1) = \{p_1\}$

Case 2 : $O_2(p_1) = \mathbb{R}^2$

Case 3 : $O_2(p_1)$ is a compact subset of \mathbb{R}^2 , star-shaped with respect to p_1 and such that $O_2(p_1) \cap L \neq \{p_1\}$ for all rays L with endpoint p_1 .

We demonstrate that case 3 cannot hold true. Suppose on the contrary that it does.

Step 10 (Convexity) : $O_2(p_1)$ is convex.

Suppose on the contrary that there exist $u, t \in O_2(p_1)$ and $w \in]t, u[$ such that $w \notin O_2(p_1)$. Consider the ray L with endpoint p_1 and containing w . By construction, we deduce $w \notin [p_1, z(L)]$. In the triangle p_1ut , either u or t is above the line $L^\perp(z)$ and therefore does not belong to $O_2(p_1)$. This argument is illustrated on figure 2.

Insert Figure 2 here

Let $p_1 = (0, 0)$. Let $(k, 0) \in O_2(p_1)$ be on the frontier of $O_2(p_1)$ and let $f : [0, k] \rightarrow \mathbb{R}_+$, defined by: $f(a) = b$ where b is the unique value of c such that (a, c) belongs to the frontier of $O_2(p_1)$. From the properties of $O_2(p_1)$, the function f is well defined. From step 10, f is concave and therefore (Rockafellar (1970)), it is left differentiable and right differentiable everywhere. Let $f'_+(a)$ and $f'_-(a)$ be the right and left derivatives of f at a .

Step 11 : $f'_+(a) = f'_-(a) \equiv f'(a)$

Without loss of generality, assume on the contrary that $f'_+(a) \leq 0 < f'_-(a)$ ⁵. Let L_- be the line with slope $f'_-(a)$ containing $(a, f(a))$. Since $O_2(p_1)$ is convex, it is included in the half plane with frontier L_- and containing p_1 . Let L be the line orthogonal to L_- containing p_1 and $w = z(L)$. Necessarily, $w = L_- \cap L$ as otherwise $(a, f(a)) \notin O_2(p_1)$ contradicting our assumption. From step 9, we deduce that $[w, (a, f(a))] \subseteq O_2(p_1)$. Since further $[w, (a, f(a))] \subseteq L_-$, we deduce that any ray L' with endpoint p_1 and intersecting $[w, (a, f(a))]$ satisfies : $z(L') = L' \cap [w, (a, f(a))]$. But this contradicts step 7. The argument is illustrated on figure 3.

Insert Figure 3 here

⁵We normalize to 0 the slope of the line orthogonal to the line generated by p_1 and $(a, f(a))$ and passing through $(a, f(a))$.

From step 11, we know that f is differentiable.

Step 12 (Geometry) : $O_2(p_1)$ is a disk centered on p_1 .

From step 7 we know that $(1, f'(a))$ is orthogonal to $(a, f(a))$ for all $a \in]0, k[$ i.e. f is solution of the differential equation:

$$a + f(a)f'(a) = 0$$

Let $F(a) = f(a)^2$. F is solution of the differential equation:

$$2a + F'(a) = 0$$

whose solutions are:

$$F(a) = C - a^2 \text{ for some constant } C$$

and therefore:

$$f(a) = \sqrt{C - a^2} \text{ for some constant } C$$

Since $f(k) = 0$, we deduce that $C = k^2$.

Step 13 : Case 3 does not hold true

From step 11 $O_2(p_1)$ is a disk centered on p_1 . Let $p_2 \notin O_2(p_1)$, $z = [p_1, p_2] \cap O_2(p_1)$ and $p'_1 \in O_2(p_1)$ such that $p_1 \in [p'_1, p_2]$. From step 5, we know that $C(p'_1, p_2) \in [p'_1, p_2]$. Let us show that $C(p'_1, p_2) = z$. Suppose not. Then either $C(p'_1, p_2) \in [p'_1, z[$ or $C(p'_1, p_2) \in]z, p_2]$. In the first case, $\|C(p'_1, p_2) - p_1\| < \|z - p_1\| = \|C(p_1, p_2) - p_1\|$ contradicting step 4. In the second case, $\|z - p'_1\| = \|C(p_1, p_2) - p'_1\| < \|C(p'_1, p_2) - p'_1\|$ contradicting also step 4.

Since $C(p'_1, p_2) \notin \{p'_1, p_2\}$, we deduce from step 11 that $O_2(p'_1)$ is a disk centered on p'_1 . Further, since $C(p'_1, p_2) = z$, the radius of the disk is $\|z - p'_1\|$. Let $p'_2 \notin O_2(p'_1) \cup [p_1, p_2]$, $v = C(p_1, p'_2)$ and $w = C(p'_1, p'_2)$ as illustrated on figure 4.

Insert Figure 4 here

Since $v = \|C(p_1, p'_2) - p'_1\| < \|C(p'_1, p'_2) - p'_1\| = w$, we contradict step 4.

At this stage, we know that for all $p_1 \in \mathbb{R}^2$, either $O_2(p_1) = \{p_1\}$ or $O_2(p_1) = \mathbb{R}^2$. The following claim shows that we can invert the quantifiers.

Step 14 (Dichotomy) : Either $O_2(p_1) = \{p_1\}$ for all $p_1 \in \mathbb{R}^2$ or $O_2(p_1) = \mathbb{R}^2$ for all $p_1 \in \mathbb{R}^2$.

Assume on the contrary that there exist $p_1, p'_1 \in \mathbb{R}^2$ such that: $O_2(p_1) = \{p_1\}$ and $O_2(p'_1) = \mathbb{R}^2$. Let p_2 be outside the disk centered on p'_1 with radius $\|p_1 - p'_1\|$. We have $C(p'_1, p_2) = p_2$ and $C(p_1, p_2) = p_1$. Since $\|p_1 - p'_1\| < \|p_2 - p'_1\|$, we contradict step 4.

The proof of the result when $n = m = 2$ is complete. When $O_2(p_1) = \{p_1\}$ for all $p_1 \in \mathbb{R}^2$, 1 is a dictator and when $O_2(p_1) = \mathbb{R}^2$ for all $p_1 \in \mathbb{R}^2$, 2 is a dictator.

4.2 The Case where $m = 2$ and $n \geq 2$

The proof is by induction on n . From the preceding section, we know that the result holds true when $n = 2$. We assume that it holds true when the number of individuals is less than $n - 1$. Let us consider the option set $O_{-1}(p_1)$ of the coalition $S = N \setminus \{1\}$ when individual 1 reports the preference p_1 .

Step 15 : Either $O_{-1}(p_1) = \{p_1\}$ or $O_{-1}(p_1) = \mathbb{R}^2$.

The proof follows with no changes the arguments from step 6 to step 13 once steps 4 and 5 have been replaced respectively by the following two properties:

$$\| C(p_1, p, p, \dots, p) - p \| \leq \| x - p \| \text{ for all } x \in O_{-1}(p_1)$$

$$C(p_1, p, p, \dots, p) \in [p_1, p]$$

The first property follows in fact from an argument similar to one implicit in step 1 applied to the mechanism $C_{-1}(p_1)$ while the second follows from efficiency. If $O_{-1}(p_1) = \{p_1\}$, then $O_{-1}(p'_1) = \{p'_1\}$ for all $p'_1 \in \mathbb{R}^2$ i.e. 1 is a dictator. Indeed, assume on the contrary that $O_{-1}(p'_1) = \mathbb{R}^2$ for some $p'_1 \in \mathbb{R}^2$. Let p_2 be outside the disk centered on p'_1 with radius $\| p_1 - p'_1 \|$. We have $C(p'_1, p_2, p_2, \dots, p_2) = p_2$ and $C(p_1, p_2, p_2, \dots, p_2) = p_1$. Since $\| p_1 - p'_1 \| < \| p_2 - p'_1 \|$, we contradict our assumption that C is strategyproof.

If $O_{-1}(p_1) \neq \{p_1\}$, then $O_{-1}(p'_1) = \mathbb{R}^2$ for all $p'_1 \in \mathbb{R}^2$. Since C is coalitional strategyproof, $C_{-1}(p_1)$ is also coalitional strategyproof. Since $R(C_{-1}(p_1)) = O_{-1}(p_1) = \mathbb{R}^2$, we deduce from the induction hypothesis that $C_{-1}(p_1)$ is dictatorial. Let $i(p_1)$ be this dictator. We now show that $i(p_1)$ is constant. Suppose that it is not constant and without loss of generality, assume that there exist $p_1, p'_1 \in \mathbb{R}^2$ such that: $i(p_1) = 2$ and $i(p'_1) = 3$. Then for all $p_2, p_3 \in \mathbb{R}^2$

$$C(p_1, p_2, p_3, \dots) = p_2 \text{ and } C(p'_1, p_2, p_3, \dots) = p_3$$

If p_2, p_3 are such that $\| p_1 - p_3 \| < \| p_1 - p_2 \|$, then the two equalities above contradict the assumption that C is strategyproof.

4.3 The Case where $m \geq 2$ and $n \geq 2$

The proof is by induction on m . From the preceding section, we know that the result is true when $m = 2$. We assume that it is true when the number of dimensions is less than $m - 1$. Let $p \in \mathbb{R}^{nm}$. If there exists an hyperplane H such that $p_i \in H$ for all $i = 1, \dots, n$, then we

deduce from the efficiency of C that $C(p) \in H$ as the convex hull of the set $\{p_1, \dots, p_n\}$ is contained in H . We deduce from the induction hypothesis that the restriction C^H of the social choice mechanism C to H^n is dictatorial. Let $i(H)$ be the dictator attached to C^H . We claim that $i(H)$ is constant. Take another hyperplane H' . If $H' \cap H \neq \emptyset$, then obviously $i(H') = i(H)$. If $H' \cap H = \emptyset$, there exist a third hyperplane H'' such that $H'' \cap H \neq \emptyset$ and $H'' \cap H' \neq \emptyset$. Then, $i(H) = i(H'') = i(H')$. Let $i(H) = 1$ for all hyperplanes H . It remains to show that 1 is a dictator.

If $n \leq m$, it follows from the argument above. If $n > m$, consider $p \in \mathbb{R}^{nm}$ and let $z = C(p)$. Let $z_n \equiv C(p_1, p_2, \dots, p_{n-1}, z)$. We claim that $z_n = z$. Indeed, if $z_n \neq z$, then since $\|C(p_1, p_2, \dots, p_{n-1}, p_n) - z\| < \|C(p_1, p_2, \dots, p_{n-1}, z) - z\|$, we would contradict our assumption that C is strategyproof. By repeating $n - m + 1$ times this argument, we obtain:

$$C(p_1, p_2, \dots, p_{m-1}, z, z, \dots, z) = z$$

Since the set $\{p_1, p_2, \dots, p_{m-1}, z\}$ is contained in an hyperplane, we conclude that $z = p_1$.

5 Related Literature

In this last section, we offer a brief and selective review of the related literature and formulate some open problems.

5.1 The Case where $m=1$

The result has been established under the assumption that there are at least two dimensions. When $m = 1$, the result does not hold. There exist coalitional strategy proofness social choice mechanisms which are not dictatorial. The exploration of the class of strategyproof mechanisms in the one dimensional setting started with the seminal paper of Moulin (1980). A complete account of this large literature can be found in Barbera (2010).

5.2 Changing the Domain and/or the Range

In this paper, a social choice mechanism has been defined as a function from \mathbb{R}^{nm} into \mathbb{R}^m . We could consider the general case of a function C from A^n into B where A and B are both subsets of \mathbb{R}^m : A and B would represent respectively the set of admissible individual ideal points and the set of feasible social alternatives. Note that if $A = \mathbb{R}^m$ and C is not surjective, in contrast to what has been assumed in our result, then B denotes simply the range of the mechanism i.e. $R(C)$. In Bordes, Le Breton and Laffond (1990), we prove that

any coalitional strategy proof onto social choice mechanism $C : \mathbb{R}^2 \rightarrow B$ where B is an equilateral triangle is dictatorial.

To the best of our knowledge, few authors have considered the general case. The unique exception is van der Stel's thesis (1993) who offers a very general and stimulating analysis of the case where $A = B$ i.e. C is a social choice mechanism from A^n onto A . He proves the following generalization of our result : if A is an open subset of \mathbb{R}^m , then any coalitional strategy proof onto social choice mechanism is dictatorial. His proof is different from ours and uses as auxilliary steps several other results.

The general study of the implications of coalitional strategy proofness for an arbitrary A seems rather intricate. Consider the case where $A = \{(0, 0), (0, 1), (1, 0), (1, 1), (\frac{1}{2}, \frac{1}{2})\}$ i.e. the set of admissible ideal points as well as the set feasible alternatives is assumed to be the set of vertices of the unit square together with its center. Consider the mechanism $C : A^n \rightarrow A$ defined as follows:

$$C(\mathbf{p}) = \begin{cases} p & \text{if } \mathbf{p} = (p, p, \dots, p) \\ (\frac{1}{2}, \frac{1}{2}) & \text{otherwise} \end{cases}$$

Since $(\frac{1}{2}, \frac{1}{2})$ is at least second best alternative for all Euclidean preferences with an ideal point in A , it is easy to check that, this mechanism is coalitional strategyproof. We could object that in this example A is not a product set. Having a product set does not help either. In the case where $A = \{(0, 0), (0, 1), (1, 0), (1, 1)\}$, the mechanism C defined by $C(\mathbf{p}) = (p_1^1, p_2^2)$ (individual 1 is a dictator on the first coordinate while individual 2 is a dictator on the second coordinate) is coalitional strategyproof. This mechanism is *decomposable*⁶ in the sense that the choice of the j^{th} coordinate only depends upon the vector $p^j = (p_1^j, p_2^j, \dots, p_n^j)$.

The finite sets A considered in the above examples are non convex but convexity of A is not necessary for the equivalence of coalitional strategyproofness and dictatorship either, as illustrated by the the case where A is the unit circle. To the best of our knowledge, the characterization of the sets A for which coalitional strategy proofness is equivalent to dictatorship is still open.

5.3 Strategyproofness and Efficiency

Can we replace, in the statement of the result, coalitional strategyproofness by (Pareto) efficiency and strategyproofness ? By definition, the answer is affirmative when $n = 2$. But

⁶A proof that decomposability follows from strategyproofness whenever the domain consists of a rich set of separable preferences appears in Le Breton and Sen (1999).

it is not when $n \geq 3$. For instance when $X = \mathbb{R}^2$ and $n = 3$, the social choice mechanism selecting the median of the coordinates of the ideal points for each of the two coordinates is surjective, strategyproof and Pareto efficient. Looking at our proof is quite instructive. While the coalitional strategyproofness property is preserved when we move from the mechanism C to the (sub)mechanism $C_{-i}(p_i)$, the Pareto efficiency property is not. The induction argument which is used in our proof breaks down.

These observations raise a new question : what are the implications of the conjunction of strategyproofness and efficiency in the Euclidean setting ? This question has been explored by Peters, van der Stel and Storcken (1992,1993a,1993b) in a series of important papers. They consider the subclass of anonymous social choice mechanisms. One striking result that Peters, van der Stel and Storcken (1992) establishes asserts that if C is a strategyproof and efficient social choice mechanism, then C is continuous. Following Moulin (1980) (for the case where $m = 1$) and Border and Jordan (1983), they define the notion of coordinatewise median scheme as follows. A collection $\{x^1, x^2, \dots, x^m\} \subseteq \mathbb{R}^m$ is a coordinate system if $\langle x^j, x^k \rangle = 0$ for all $j, k = 1, \dots, m$ with $j \neq k$ i.e. $\{x^1, x^2, \dots, x^m\}$ is an orthogonal basis of \mathbb{R}^m . Let $k \in \mathbb{N}$ so that $k+n$ is odd. A social choice mechanism C is a *coordinatewise median social choice mechanism* with k constant points (phantom points) if there exists a coordinate system and points $c_1, c_2, \dots, c_k \in \{\mathbb{R} \cup \{-\infty, \infty\}\}^m$ such that:

$$C^j(p) = Med(p_1^j, p_2^j, \dots, p_n^j, c_1^j, c_2^j, \dots, c_k^j)$$

where Med denotes the median of the subsequent real numbers and all coordinates are expressed with respect to the given coordinate system. They show that if $m \geq 2$ and n is even, or if $n \geq 3$, then there does not exist surjective social choice mechanism which are anonymous, efficient and strategyproof. However when $m = 2$ and n is odd, the class of anonymous, efficient and strategyproof social choice mechanisms is non empty. More precisely, the class of anonymous, efficient and strategyproof social choice mechanisms coincides exactly with the class of coordinatewise median social choice mechanism without constant points. This result was established also by Kim and Roush (1984) with the additional assumption that C is continuous. But, as already pointed out by Peters, van der Stel and Storcken, this property is implied by the others and is therefore redundant.

5.4 Strategyproofness

Coalitional strategyproofness is often strictly more demanding than strategyproofness. It is therefore natural to wonder what is the class of strategyproof social choice mechanisms

when $X = \mathbb{R}^m$. The coordinatewise median social choice mechanisms are strategyproof. More generally, we can construct componentwise social choice mechanisms by constructing separately a social choice mechanism for each of the m components where the choice of the j^{th} social coordinate only depends upon the vector $p^j = (p_1^j, p_2^j, \dots, p_n^j)$. Such mechanisms have been called *decomposable* as the initial problem has been decomposed into m one dimensional problems. Strategyproofness follows from the fact that Euclidean preferences are separable i.e. preferences over one coordinate do not depend upon what has been decided upon the other coordinates. Therefore, existence of non dictatorial social mechanisms is not an issue here. It is legitimate to wonder what is the exact class of strategyproof social choice mechanisms.

This a difficult question which has been adressed by Laffond (1980) in his thesis and also by Kim and Roush (1984). Kim and Roush focus on the case where $X = \mathbb{R}^2$. They prove that a social choice mechanism is continuous and anonymous (but not necessarily surjective) iff it is a coordinatewise median social choice mechanism with $n + 1$ constant points. Peters, van der Stel and Storcken (1993b) proves that if C is strategyproof then C is continuous iff $R(C)$ is convex. They further point out that if C is surjective then C is strategy proof and anonymous iff it is a coordinatewise median social choice mechanism with $n - 1$ constant points

Laffond's work (1980) considers the case where $n = 2$ but m is arbitrary. He focuses on the class of anonymous, continuous, surjective and strategyproof social choice mechanisms. As already pointed out, to conduct his analysis, he also invented the technique of option sets. His analysis consists in a gradual exploration of the properties of the sets $O_2(p_1)$ and $O_1(p_2)$. First, he demonstrates that if C is surjective, continuous and strategyproof, the option sets are closed and convex subsets of \mathbb{R}^m . This implies that $C(p_1, p_2)$ is the projection of p_1 on $O_1(p_2)$ and the projection of p_2 on $O_2(p_1)$. Of course some consistency condition is needed since the two options sets cannot be constructed independently of each other. The main part of his work consists in sorting out the implications of this consistency. We have no space here to go through all his lengthy analysis. We just sketch some of his main ideas. To show what kind of mechanisms will appear out of Laffond's exploration, consider, for the sake of illustration⁷, the case where $C(p_1, p_2)$ is the projection of p_2 on $p_1 + K_2$ where K_2 is the convex cone $\{x \in \mathbb{R}^2 : x_1 \leq 0, x_2 \leq 0 \text{ and } x_2 \leq x_1\}$. By construction, $O_2(p_1) = p_1 + K_2$ and C is not manipulable by 2. It is easy to show that C is not manipulable either. Further, it can be verified that $O_1(p_2) = p_2 + K_1$ where $K_1 = -K_2^\perp$ where $K_2^\perp \equiv \{x \in \mathbb{R}^2 : \langle x, y \rangle \leq 0 \text{ for all } y \in K_2\}$. K_2^\perp is called the *polar of the cone* K_2 . The construction

⁷This example appears in Border and Jordan (1983).

is illustrated on figure 5.

Insert Figure 5 here

Another illustration is the case where individual 1 is a dictator on the first coordinate and individual 2 is a dictator on the second coordinate. In such case, K_2 is the vertical axis and K_1 is the horizontal axis. In the first example, K_1 has a non empty interior while its is empty in the second one. These two examples violates anonymity. Anonymity prevails iff $K_2^\perp = -K_1$. Such cones are called *self-polar* (or self dual cones). Let K be an arbitrary closed and convex self-polar cone with an non empty interior. As shown by Laffond, the social choice mechanism where equivalently the social outcome is the projection of p_1 on $p_2 + K$ or the projection of p_2 of 2 on $p_1 + K$ is strategyproof, anonymous and continuous. This construction leads to a large family of anonymous, surjective, continuous and strategyproof social choice mechanism. What mechanisms do we find in that family ?

First, and not surprisingly (from what precedes), we find the class of coordinatewise median social choice mechanism with one constant point at infinity. The self-polar cone attached to any such mechanism is (up to a rotation) the positive orthant \mathbb{R}_+^m . When $m = 2$, \mathbb{R}_+^2 is the unique (up to a rotation) self-polar cone of \mathbb{R}^2 . When $m = 2$, any social choice mechanism constructed along these lines is decomposable. However, when $m \geq 3$, this *is not* true anymore. This follows from the fact that when $m \geq 3$, the family of self-polar cones is much richer. The analysis of these objects is a well defined area in mathematics (see e.g. Barker and Foran (1976), Lochum (1984)) and one merit of Laffond's construction is to point out this connection.

The above construction raises the following question : What is the specificity of the class of anonymous, surjective, continuous and strategyproof social choice mechanisms constructed by Laffond through self-polar cones ? Laffond shows that, to answer the above question, we have to examine the asymptotic cone⁸ of the option set $O(p)$. In particular, he first shows that if C is an anonymous, surjective, continuous and strategyproof social choice mechanism, then $O_1 = O_2 \equiv O$ where O is a correspondence with closed and convex values such that $K(O(p))$ is independent of p i.e. $K(O(p)) = K(O(p')) \equiv K$ for all $p, p' \in \mathbb{R}^m$. He further shows that if K has a non empty interior, then C belongs to the class which has just been constructed i.e. $O(p) = p + K$ and K is self-polar.

⁸The asymptotic cone $K(A)$ of an arbitrary closed and convex subset A of \mathbb{R}^m is defined as follows:
 $K(A) = \{x \in \mathbb{R}^m : \forall \lambda \geq 0, \forall y \in A, x + \lambda y \in A\}$

5.5 Superdomains

The set of Euclidean preferences on \mathbb{R}^m is in one to one correspondence with \mathbb{R}^m . Many authors have explored the questions examined in this note for sets of preferences who are supersets of the set of Euclidean preferences. One important such a superdomain is the set of *diagonal (separable) quadratic* preferences. They are described by utility functions u such that:

$$u(x) = - \sum_{j=1}^m \alpha^j (x^j - p^j)^2 \text{ where } \alpha = (\alpha^j)_{1 \leq j \leq m} \in \mathbb{R}_{++}^m$$

Another (larger) superdomain is the all set of *quadratic* preferences described by utility functions u such that:

$$u(x) = - \sum_{k=1}^m \sum_{j=1}^m \alpha^{kj} (x^j - p^j)^2 \text{ where } \alpha = (\alpha^{kj})_{1 \leq k, j \leq m} \text{ is a symmetric positive definite matrix}$$

Border and Jordan (1983) 's seminal contribution contains many important results on these two domains. They demonstrate that a surjective social choice mechanism C over the domain of diagonal quadratic preference is strategyproof iff it is decomposable i.e. if there exist m surjective and strategyproof social choice mechanisms C^j such that:

$$C(\mathbf{p}, \alpha) = (C(p^1), C(p^2), \dots, C(p^m))$$

Among other things, decomposability obliges to abandon the idea of using the information on preferences which is contained in the vectors $\alpha_i \in \mathbb{R}_{++}^m$ and therefore to limit the information on preferences to the peaks of the individual preferences⁹. This decomposability result implies here a property which has been called in the literature a *tops only property*¹⁰. In many other social environments settings, it is also an implication of strategyproofness.

They prove that any surjective and strategyproof social choice mechanism over the set of quadratic preferences is dictatorial. Their result has been generalized by Zhou (1991) who weakens the surjectivity assumption to a very weak range condition.

The superdomains discussed above are *parametric*. We could consider classes of preferences which are not described by a finite set of parameters. For example we could consider the class of preferences on \mathbb{R}^m represented bu utility functions u such that:

$$u(x) = \sum_{j=1}^m v^j(x^j)$$

⁹Decomposability implies more restrictions.

¹⁰On this, see Weymark (1999).

where each $v^j : \mathbb{R} \rightarrow \mathbb{R}$ has a unique maximizer from which it decreases monotonically in either direction. This domain has been explored by many authors including among others Barbera, Gul and Stachetti (1993). It is important to note that if it can be shown that strategyproofness implies the top only property, then we are back to the parametric domains considered earlier. Barbera (2010) offers a detailed exposition of this area of research.

Finally, we could also consider the questions explored in this note for a class of preferences generated by a distance different from the Euclidean distance. This question is explored extensively in van der Stel who considers some other norms. In location problems, the set A is assumed to be a closed and connected subset of \mathbb{R}^m and for all $p, q \in A$, the distance between $d(p, q)$ is the shortest distance from p to q (and q to p). Schummer and Vohra (2002) have examined the implications of surjectivity and strategyproofness in the case where A is the union of a finite number of closed curves of finite length.

6 References

Barbera, S. (2010) Strategyproof Social Choice, Chapter 25 in *Handbook of Social Choice and Welfare*, Vol 2, 731-831.

Barbera, S., Gul, F. and E. Stacchetti (1993) "Generalized Median Voter Schemes and Committees", *Journal of Economic Theory*, 61, 262-289.

Barbera, S., Masso J. and A. Neme (1997) "Voting Under Constraints", *Journal of Economic Theory*, 76, 298-321.

Barbera, S., Masso, J. and S. Serizawa (1998) "Strategy-Proof Voting on Compact Ranges", *Games and Economic Behavior*, 25, 272-291.

Barbera, S. and Peleg (1990) "Strategy-Proof Voting Schemes with Continuous Preferences", *Social Choice and Welfare*, 7, 31-38.

Barker, G.P. and J. Foran (1976) "Self-Dual Cones in Euclidean Spaces", *Linear Algebra and its Applications*, 13, 147-155.

Berga, D. and S. Serizawa (2000) "Maximal Domains for Strategy-Proof Rules with One Public Good", *Journal of Economic Theory*, 90, 39-61

Border, K. and J. Jordan (1983) "Straightforward Elections, Unanimity and Phantom Voters", *Review of Economic Studies*, 50, 153-170.

Bordes, G., Le Breton, M. and G. Laffond (1990) "Strategy-Proofness Issues in Some Economic and Political Domains", Mimeo.

Kim, K.H. and F.W. Roush (1984) "Nonmanipulability in Two Dimensions", *Mathematical Social Sciences*, 8, 29-43.

- Iochum, B. (1984) *Cônes Autoplaires et Algèbres de Jordan*, Lecture Notes in Mathematics N° 1049, Springer-Verlag.
- Laffond, G. (1980) *Révélation des Préférences et Utilités Unimodales*, Thesis, Paris.
- Laffont, J.J. (1987) "Revelation of Preferences" in *The New Palgrave : A Dictionary of Economics*, J. Eatwell, M. Milgate and P. Newman (Eds), First Edition, Palgrave Mc Milan.
- Le Breton, M. and A. Sen (1999) "Separable Preferences, Strategyproofness and Decomposability", *Econometrica*, 67, 605-628.
- Moulin, H. (1980) "On Strategy-Proofness and Single Peakdedness", *Public Choice*, 35, 437-455.
- Peters, H., van der Stel, J. and T. Storcken (1992) "Pareto Optimality, Anonymity and Strategy-Proofness in Location Problems", *International Journal of Game Theory*, 21, 221-235.
- Peters, H., van der Stel, J. and T. Storcken (1993a) "Generalized Median Solutions, Strategy-Proofness and Strictly Convex Norms", *Zeitschrift Für Operations Research*, 38, 19-53.
- Peters, H., van der Stel, J. and T. Storcken (1993b) "Range Convexity, Continuity and Strategy-Proofness of Voting Schemes", *Zeitschrift Für Operations Research*, 38, 213-229.
- Schummer, J. and R.V. Vohra (2002) "Strategy-Proof Location on a Network", *Journal of Economic Theory*, 104, 405-428.
- van der Stel, J. (1993) *Strategy-Proof Voting Schemes on Euclidean Spaces*, Thesis, Maastricht.
- Rockafellar, R.T. (1970) *Convex Analysis*, Princeton University Press, New Jersey.
- Weymark, J.A. (1999) "Decomposable Strategy-Proof Social Choice Functions", *Japanese Economic Review*, 50, 343-355.
- Zhou, L. (1991) "Impossibility of Strategy-Proof Mechanisms in Economies with Pure Public Goods", *Review of Economic Studies*, 58, 107-119.