

Kinetic models for topological nearest-neighbor interactions

Adrien Blanchet and Pierre Degond



Kinetic models for topological nearest-neighbor interactions

March 16, 2017

Adrien Blanchet

IAST/TSE, Université Toulouse Capitole – 21 Allée de Brienne 31000
Toulouse, France
email: Adrien.Blanchet@ut-capitole.fr

Pierre Degond

Department of Mathematics, Imperial College London, London SW7 2AZ,
United Kingdom
email: p.degond@imperial.ac.uk

Abstract

We consider systems of agents interacting through topological interactions. These have been shown to play an important part in animal and human behavior. Precisely, the system consists of a finite number of particles characterized by their positions and velocities. At random times a randomly chosen particle, the follower adopts the velocity of its closest neighbor, the leader. We study the limit of a system size going to infinity and, under the assumption of propagation of chaos, show that the limit kinetic equation is a non-standard spatial diffusion equation for the particle distribution function. We also study the case wherein the particles interact with their K closest neighbors and show that the corresponding kinetic equation is the same. Finally, we prove that these models can be seen as a singular limit of the smooth rank-based model previously studied in [10]. The proofs are based on a combinatorial interpretation of the rank as well as some concentration of measure arguments.

Keywords: rank-based interaction, spatial diffusion equation, continuity equation, concentration of measure

1 Introduction

In the literature on animal behavior including fish [3], birds [38] and even pedestrians [32], interactions between individuals are often assumed to be strongly dependent on their relative distance. However it has recently been demonstrated

that individuals in bird flocks interact with their nearest neighbors irrespective of their distance [5, 18]. More precisely, the authors in [5] claim that each bird interacts with between six to eight of its closest neighbors. The authors coined the term of “topological interaction” to refer to such interaction mechanisms and “topological distance” to refer to how many other individuals were closer. Even though the reality of this topological interaction has been debated [24], it now seems to receive consensus following reports that self-propelled particle models based on topological interactions successfully reproduce the observed experimental features [11, 14, 28].

The understanding that birds interact through topological rather than metric distance has generated an intense literature. Topological interactions have been introduced to the modeling of many natural phenomena, from birds [33] to pedestrians [36]. Mixed metric-topologic interactions have also been proposed [41, 44]. Proof of flocking under topological interaction has been given in [31, 39, 46], while speed to consensus has been shown to depend on the number of interacting neighbors in [45]. Similarly, interactions depending of the behavior of the closest neighbor are probably at play in human interactions such as portfolio theory [6, 25, 35], competition between coworkers within a firm, risk-taking among traders or aggressive behavior to reach a sexual partner, see for example [22, 34]. Rank-based dynamics bears similarities with rearrangement (see e.g. [13] and references therein).

One of the striking features of topological compared to metric interactions is their scale-invariance property. Indeed, irrespective of the bird concentration within the flock, [5] proved that the interaction features remain unchanged. In human cognition models, it is also more relevant to restrict interactions to the closest neighbors, as the attention of a subject is intrinsically directed to only a few people around him/her [42]. Interactions with close neighbors do not preclude interactions at a longer range, as interactions spread with the conscious or unconscious signals sent by the subjects in response to these interactions. However, issues such as quantify the propagation speed of information sent via topological interactions are poorly understood so far, in particular in the presence of a large number of subjects. This calls for a large-scale theory of topological interactions, or in other words, for the development of meso or macroscopic models of particle systems connected through topological interactions. The aim of this article is specifically to derive a macroscopic model for a large population of particles interacting through topological interactions, starting from a simple microscopic model. To the best of our knowledge, the present work and its predecessor [10] are the first to develop a rigorous coarse-graining of topological interactions, with the exception of [31] which tackled a similar question but for a different kind of interaction, closer to mean-field type interactions.

More precisely, we will consider a system with interacting mobile agents. At Poisson random times a given agent selects a partner to interact with according to a probability rule which depends on the proximity rank of the partner. The interaction rule is then very simple: the agent changes its velocity to align with that of its partner. The goal of the present work is, by letting the number of agents tend to infinity, to derive an equation for the probability distribution

of the agents in phase space (positions, velocities). Thanks to the choice of this simple interaction rule, inspired from earlier work [15, 16], we can concentrate on the mathematical aspects of this derivation. In previous work by the authors [10], the probability rule solely depended on the proximity rank normalized by the total number of interacting partners (or equivalently, on a proximity rank expressed as a percentage). This rule made the number of potential interacting partners tend to infinity as the number of agents also did so. We will refer to this rule as the "smooth rank-based dynamics". By contrast, in the present work, we will concentrate on the case where there is only one interaction partner (the nearest one) or a finite number of them (the K nearest ones), even in the limit of the number of agents tending to infinity. We will refer to these dynamics as the "nearest-neighbor" or " K -nearest-neighbor" dynamics.

In this paper, we show (under the Propagation of Chaos assumption) that the kinetic equation resulting from the nearest-neighbor or K -nearest neighbor dynamics is a nonlinear spatial diffusion equation for the particle distribution function in phase space (position, velocity). This equation has a non-classical feature as it involves a spatial anti-diffusion of the density (which is a velocity integral of the distribution function). We will show that this term results from the constraint that the density must satisfy the continuity equation, a constraint resulting from the preservation of the number of particles in the course of an interaction. By contrast, in the previous work [10] relative to the smooth rank-based dynamics, we showed (also under the Propagation of Chaos assumption), that the resulting kinetic equation involved a spatially non-local integral equation and that the continuity equation for the density was also satisfied. In the present paper, we also show that we can pass from the non-local integral equation issued from the smooth rank-based dynamics to the nonlinear spatial diffusion equation for the nearest-neighbor dynamics by a process involving a singular concentration of the kernel of the integral equation. This provides a vision of the nearest-neighbor dynamics as a singular limit of the previously studied smooth rank-based dynamics.

Rank-based dynamics (either smooth or nearest-neighbor) are natural from a mathematical point of view. The proximity rank includes information about the most immediate interaction partners of a given particle. Although the rank is a highly non-linear function of the particle positions and is subject to jumps when two particles cross, it has robust properties such as invariance by permutation of the particle numbers, and has combinatorial interpretation: the probability for an agent to have a rank k with respect to agent i is equal to the probability of having $k - 1$ agents between them. Our results strongly rely on this interpretation of the rank, together with some concentration of measure arguments. Rank-based dynamics also exhibit a certain universality, as their kinetic model does not depend on the (finite) number of interacting agents. The kinetic equation that we derive in this work leads to many questions. This equation is a non-standard diffusion equation with mainly unknown properties, from the perspective of well-posedness, large-time behavior, regularity, etc. We believe that these questions open fascinating new directions of research in kinetic theory.

Kinetic models of flocking or swarming behavior have been widely inves-

tigated in the context of metric interactions. The literature is vast and it is virtually impossible to be exhaustive. Below is a sample of major publications on this topic. Derivation of kinetic models from underlying particle models have been established in [12, 21, 30, 43]. Flocking behavior and pattern formation has been investigated in [1, 9, 17, 29, 30, 40]. Equilibria and phase transitions in kinetic flocking models have been studied in [7, 19, 20]. Passage from kinetic to hydrodynamic descriptions of flocking has been investigated in [8, 19, 20, 21, 23, 26]. Numerical simulation methods have been put forward in [2, 27].

The organization of the paper is as follows. Section 2 is dedicated to the presentation of the models and our main results, as well as a detailed discussion of them. Section 3 covers the nearest-neighbor case and the derivation of the kinetic model in this case. Section 4 extends these results to the case where the particles interact with their K closest neighbors. Section 5 develops the proof that the kinetic model of the nearest-neighbor (or K -nearest-neighbor) interaction is the limit of the kinetic smooth rank-based interaction model of [10]. A conclusion to this article is given in Section 6.

2 Models and main results

2.1 General framework

Consider a set of N particles. The particle i is characterized by its position $x_i \in \mathbb{R}^n$ and its velocity $v_i \in \mathbb{R}^n$ where $n \geq 1$ is both the spatial and velocity dimension. Let $(\pi_{ij}^N)_{(i,j) \in \{1, \dots, N\}^2}$ be a stochastic matrix, *i.e.* for all $(i, j) \in \{1, \dots, N\}^2$, $\pi_{ij}^N \in [0, 1]$ and for all $i \in \{1, \dots, N\}$, $\sum_{j=1}^N \pi_{ij}^N = 1$. The particles $\{(x_1(t), v_1(t)), \dots, (x_N(t), v_N(t))\}$ are subject to the following dynamics

- The dynamics is a succession of free-flights and collisions.
- During free-flight, particles follow straight trajectories

$$\begin{cases} \dot{x}_i = v_i, \\ \dot{v}_i = 0. \end{cases}$$

- At Poisson random times with a rate equals to $N \lambda(N)$, particles undergo the following collisions process: pick a particle i in $\{1, \dots, N\}$ with uniform probability $1/N$; then pick a collision partner j with probability π_{ij}^N and perform the collision:

$$\begin{cases} (x_i, x_j) \text{ remains unchanged,} \\ (v_i, v_j) \text{ is changed into } (v_j, v_j). \end{cases}$$

We will assume that π_{ij}^N is a function of the particle positions (x_1, \dots, x_N) , *i.e.* $\pi_{ij}^N = \pi_{ij}^N(x_1, \dots, x_N)$ and is permutation invariant, *i.e.* for any permutation

$\sigma \in \mathfrak{S}_N$ where \mathfrak{S}_N denotes the set of permutations of $\{1, \dots, N\}$, we have

$$\pi_{\sigma(i)\sigma(j)}^N(x_{\sigma(1)}, \dots, x_{\sigma(N)}) = \pi_{ij}^N(x_1, \dots, x_N). \quad (1)$$

The function $\lambda(N)$ is an appropriate scaling factor which will be defined and discussed below.

To simplify the notation, when no confusion is possible, we will denote $\mathbf{x} := (x_1, \dots, x_N)$, $\mathbf{v} := (v_1, \dots, v_N)$, $Z_i := (x_i, v_i)$, $\mathbf{Z} := (Z_1, \dots, Z_N)$ and $d\mathbf{Z} := dx_1 dv_1 \dots dx_N dv_N$. We will also use $f(d\mathbf{Z})$ instead of $f(\mathbf{Z}) dx$.

The system will be described through the master equation which provides the dynamics of the N -particles distribution $f^{(N)}(Z_1, \dots, Z_N)$, i.e. the joint probability of particles 1 to N to be at location (in phase space) Z_1 to Z_N . As shown in [10], thanks to (1) if $f^{(N)}(t)$ is permutation invariant at time $t = 0$ then it is permutation invariant for all times. We study the limit of this dynamics when the number of particles goes to ∞ . First define the k -th marginal ($k \in \{1, \dots, N\}$) as

$$f_N^{(k)}(Z_1, \dots, Z_N, t) := \int f^{(N)}(\mathbf{Z}) dZ_{k+1} \dots dZ_N.$$

We assume that ‘‘Propagation of Chaos’’ holds true *i.e.* $\forall \mathbf{Z} \in \mathbb{R}^{2nN}, \forall t \in [0, \infty)$:

$$f^{(N)}(Z_1, \dots, Z_N, t) = \prod_{\ell=1}^N f_N^{(1)}(Z_\ell, t) + \text{negligible terms as } N \rightarrow \infty. \quad (2)$$

Assuming that $f_N^{(1)} \rightarrow f$ in the limit $N \rightarrow \infty$, we aim to derive the equation satisfied by f .

In this article we are interested in situations in which a population of agents interact not based on their metric distance but on their topological distance. To do so we introduce the following definition:

Definition 1 (Rank). *Consider N particles located at x_1, \dots, x_N . Consider the i -th particle and order the list $(|x_j - x_i|)_{j=1, \dots, N, j \neq i}$ by increasing order and denote by $r^N(i, j) \in \{1, \dots, N-1\}$ the position of the j -th item in this list. If two indices j and j' are such that $|x_j - x_i| = |x_{j'} - x_i|$, then we choose arbitrarily an ordering between these two numbers. We define $r^N(i, i) = 0$. Now, we define the rank of j with respect to i as:*

$$R^N(i, j) = \frac{r^N(i, j)}{N-1} \in \bigcup_{k=1}^{N-1} \left\{ \frac{k}{N-1} \right\}.$$

Remark 1. *The notation r and R have been interchanged compared with [10].*

We assume that the interaction probabilities π_{ij}^N depend on the particle positions (x_1, \dots, x_N) only through the rank of j with respect to i , i.e. $\pi_{ij}^N = \Pi(r^N(i, j))$ where the function Π will take different forms according to the chosen model. Since the rank of j with respect to i is an intrinsic property of the positions of the particles and does not depend on how they are numbered, we have the following properties of the rank:

Remark 2. Let (x_1, \dots, x_N) be a set on N particles.

(i) The rank $r^N(i, j)$, and hence π_{ij}^N , is a function of (x_1, \dots, x_N) , i.e.

$$r^N(i, j) = r^N(i, j)(x_1, \dots, x_N) .$$

More precisely, we consider the rank $r^N(i, j)$ as a function of $L^\infty(\mathbb{R}^{nN})$.

(ii) The rank is permutation invariant, i.e. for any permutation $\sigma \in \mathfrak{S}_N$ where \mathfrak{S}_N denotes the set of permutations of $\{1, \dots, N\}$, we have

$$r^N(\sigma(i), \sigma(j))(x_{\sigma(1)}, \dots, x_{\sigma(N)}) = r^N(i, j)(x_1, \dots, x_N) .$$

We will be considering three different type of collision dynamics which we will describe in the following three sections.

2.2 Smooth rank-based dynamics

This case has been studied in [10]. We recall it here for reference and further comparisons with the other models introduced below. We introduce a function $K: R \in [0, 1] \mapsto K(R) \in [0, \infty)$ such that

$$\int_0^1 K(R) \, dR = 1 .$$

We define

$$K^N(R) = \frac{K(R)}{\sum_{k=1}^{N-1} K\left(\frac{k}{N-1}\right)} ,$$

in order to have for any $i \in \{1, \dots, N\}$:

$$\sum_{\substack{j=1 \\ j \neq i}}^N K^N(R^N(i, j)) = \sum_{k=1}^{N-1} K^N\left(\frac{k}{N-1}\right) = 1 .$$

In this way, for any $i \in \{1, \dots, N\}$, the collection $(\pi_{ij}^N)_{j=1, j \neq i}^N$, where

$$\pi_{ij}^N = K^N(R^N(i, j)) ,$$

defines a discrete probability measure on the set $\{j \in \{1, \dots, N\}, j \neq i\}$. The collision rule is then as follows

Defintion 2 (Smooth rank-based dynamics). *Pick a particle i in $\{1, \dots, N\}$ with uniform probability $1/N$; then pick a particle $j \neq i$ with probability $\pi_{ij}^N = K^N(R^N(i, j))$ and perform the collision:*

$$\begin{cases} (x_i, x_j) \text{ remains unchanged,} \\ (v_i, v_j) \text{ is changed into } (v_j, v_j). \end{cases}$$

Assuming that $f_N^{(1)} \rightarrow f$ and $\rho_N^{(1)} := \int f_N^{(1)} dv \rightarrow \rho = \int f dv$, as $N \rightarrow \infty$, and assuming $\lambda(N) = 1$, we prove in [10] that f is a solution of the kinetic equation:

$$\frac{\partial f}{\partial t}(x, v) + v \cdot \nabla_x f(x, v) = \rho(x) \int f(x', v) K(M_\rho(x, |x' - x|)) dx' - f(x, v), \quad (3)$$

where M_ρ is the partial mass of ρ and is defined by

$$M_\rho(x, s) = \int_{x' \in |x - x'| \leq s} \rho(x') dx'. \quad (4)$$

2.3 Nearest neighbor dynamics

In this article we first consider the case where the collision takes place with the closest neighbor *i.e.*

$$\pi_{ij}^N = \delta_1(r^N(i, j))$$

where δ_x stands for the Dirac delta centered at x .

We will prove the following: Assume Propagation of Chaos (2) and assume a specific form for $\lambda(N)$ to be made precise below. Then, if $f_N^{(1)} \rightarrow f$ and $\rho_N^{(1)} := \int f_N^{(1)} dv \rightarrow \rho = \int f dv$ as $N \rightarrow \infty$, f and ρ satisfy

$$\partial_t f(x, v) + v \cdot \nabla_x f(x, v) = Q[f](x, v), \quad (5)$$

where the collision operator $Q[f]$ is defined by

$$Q[f] := \frac{1}{\rho^{\frac{d}{2}+1}} \left[\Delta_x f - \frac{f}{\rho} \Delta_x \rho \right]. \quad (6)$$

Note that we do not intend to give a fully rigorous derivation of this model but of course some assumptions have to be made to ensure that ρ for instance remains positive in the considered domain. In the same spirit the topology in which the convergence takes place will be omitted here and discussed in future works.

2.4 K -nearest neighbor dynamics

Let K be a fixed integer in $\{1, \dots, N\}$ and let $(\alpha_k)_{k \in \{1, \dots, K\}}$ be a given sequence such that $\sum_{k=1}^K \alpha_k = 1$. We extend this sequence for k in $\{1, \dots, N\}$ by taking $\alpha_k = 0$ for all $k > K$. We consider a collision rule by which particle i adopts the j th-particle velocity with probability α_k if particle j is particle i 's k -th nearest neighbor, *i.e.* the probability π_{ij} is given by $\pi_{ij} = \sum_{k=1}^N \alpha_k \delta_k(r^N(i, j))$. Section 2.3 on nearest neighbor interaction was dedicated to the case $\alpha_1 = 1$ and $\alpha_k \equiv 0$ for all $k > 1$, *i.e.* $K = 1$. The rule considered here is a generalization to the K nearest neighbors *i.e.* :

$$\pi_{ij}^N(\mathbf{x}) = \sum_{k=1}^N \alpha_k \delta_k(r^N(i, j)(\mathbf{x})). \quad (7)$$

Under the propagation of chaos (2) and upon choosing a convenient scaling function $\lambda(N)$, we will prove that f satisfies the same equation (5) as before, with the same collision operator (6).

2.5 Discussion

As we can see from Sections 2.3 and 2.4, the kinetic models derived from the nearest neighbor interaction or the K -nearest neighbor interactions are the same. On the other hand, these models differ quite significantly from that obtained from the smooth ranked-based dynamics of [10] as recalled in Section 2.2. However, we show in Section 5 that the former are limits of the latter when the interaction kernel K concentrates (with a convenient scaling) near zero. In this limit, since only the closest neighbors interact, and these closest neighbors are likely to be spatially close (especially when the density is large) the spatially non-local integral operator appearing in (3) converges to the diffusion operator (6). Note that this diffusion is multiplied by an inverse power of the density. This is easily understood as, when the density is small, the particles are very far apart, resulting in spatial communications between the particles over larger distances, and eventually, into a larger diffusion coefficient. This interpretation is reinforced by the fact that the inverse power of the density depends on the dimension, in the same way as the scaling between the inter-particle distance and the density depends on the dimension.

In [10], we noted that any solution of the smooth rank-based dynamic kinetic model (3) satisfies the mass conservation equation

$$\partial_t \rho + \nabla_x \cdot (\rho u) = 0,$$

with $\rho = \int f \, dv$ and $\rho u = \int f v \, dv$. The nearest-neighbor kinetic model (5) also satisfies the mass conservation equation. To see this, it is enough to show that $\int Q(f) \, dv = 0$. But we easily check that it is the case. Indeed:

$$\begin{aligned} \int Q(f) \, dv &= \frac{1}{\rho^{\frac{2}{d}+1}} \left[\Delta_x \left(\int f \, dv \right) - \frac{1}{\rho} \left(\int f \, dv \right) \Delta_x \rho \right] \\ &= \frac{1}{\rho^{\frac{2}{d}+1}} [\Delta_x \rho - \Delta_x \rho] = 0. \end{aligned}$$

The collision operator (6) has the form of a spatial diffusion of f but with an anti-diffusion in ρ . In fact, this anti-diffusion is exactly the term that needs to be added to turn a pure spatial diffusion $\rho^{-\frac{2}{d}-1} \Delta_x f$ into an operator that conserves mass i.e. that satisfies $\int Q(f) \, dv = 0$. The well-posedness theory of (5) is still open but from this remark we can conjecture that the model is well-posed. Indeed, apart from a mass-carrying component, the equation is a spatial diffusion. And the mass carrying component satisfies a continuity equation. So, it seems that the model couples two components each of which solves a well-posed equation. Of course, the coupling is non-trivial and this may result into a lack of well-posedness. This issue will be dealt with in future work.

3 Nearest neighbor interaction

3.1 Master equation and propagation of chaos

3.1.1 Master equation

As recalled in [10], when the collisions occur at Poisson times with rate $N\lambda(N)$, the master equation for the N -particle distribution function $f^{(N)}$ in weak form is, for all test function $\phi^N : \mathbf{Z} \mapsto \phi^N(\mathbf{Z})$, given by:

$$\begin{aligned} \int \partial_t f^{(N)}(\mathbf{Z}) \phi^N(\mathbf{Z}) d\mathbf{Z} &= \sum_{i=1}^N \int f^{(N)}(\mathbf{Z}) (v_i \cdot \nabla_{x_i}) \phi^N(\mathbf{Z}) d\mathbf{Z} \\ &+ \lambda(N) \sum_{\substack{i,j=1 \\ j \neq i}}^N \iint \delta_1[r^N(i,j)(\mathbf{x})] [\phi^N(Z_1, \dots, x_i, v_j, \dots, x_j, v_j, \dots, Z_N) - \phi^N(\mathbf{Z})] \\ &f^{(N)}(\mathbf{Z}) d\mathbf{Z}, \quad (8) \end{aligned}$$

where δ_1 denotes the Dirac Delta centered at 1.

3.1.2 Propagation of chaos and first marginal

The following proposition provides the equation for the first marginal $f_N^{(1)}$ under the Propagation of Chaos Assumption (2). This equation is given in weak form by using a test function $\phi(Z_1)$ only depending on the first coordinate Z_1 in Eq. (8) and inserting (2) into (8). In all this section, we drop the drift term (the first term at the right-hand side of (8)) as its treatment is classical. This leads to the

Proposition 3 (First marginal equation with propagation of chaos). *Under the propagation of chaos assumption (2), for all test function $\phi(Z_1)$ and dropping the drift term as well as the negligible terms when $N \rightarrow \infty$ in (2), we have:*

$$\begin{aligned} &\frac{1}{\lambda(N)(N-1)} \int \partial_t f_N^{(1)}(Z_1) \phi(Z_1) dZ_1 \\ &= \int [\phi(x_1, v_2) - \phi(Z_1)] \left(1 - M_{\rho_N^{(1)}}(x_1, |x_2 - x_1|)\right)^{N-2} f_N^{(1)}(dZ_1) f_N^{(1)}(dZ_2), \quad (9) \end{aligned}$$

with M_ρ given by (4).

Proof. As pointed out above, we drop the drift term for simplicity as this term can be handled with classical methods. Taking $\phi^N(\mathbf{Z}) = \phi(Z_1)$ as a test function in the master equation (8) and using the permutation invariance as well as a

straightforward combinatorial argument shows that

$$\begin{aligned}
& \frac{1}{\lambda(N)(N-1)} \int \partial_t f^{(N)}(\mathbf{Z}) \phi^N(Z_1) d\mathbf{Z} \\
&= \int \delta_1[r^N(1,2)(\mathbf{x})] [\phi(x_1, v_2) - \phi(Z_1)] f^{(N)}(\mathbf{Z}) d\mathbf{Z} \\
&+ \int \delta_1[r^N(2,1)(\mathbf{x})] [\phi(Z_1) - \phi(Z_1)] f^{(N)}(d\mathbf{Z}) \\
&+ (N-2) \int \delta_1[r^N(2,3)(\mathbf{x})] [\phi(Z_1) - \phi(Z_1)] f^{(N)}(d\mathbf{Z}).
\end{aligned}$$

The last two terms obviously vanish.

Using the propagation of chaos assumption (2) we obtain

$$\begin{aligned}
& \frac{1}{\lambda(N)(N-1)} \int \partial_t f_N^{(1)}(\mathbf{Z}) \phi(Z_1) d\mathbf{Z} \\
&= \int [\phi(x_1, v_2) - \phi(Z_1)] \left\{ \int \delta_1[r^N(1,2)(\mathbf{x})] \prod_{\ell=3}^N d\rho_N^{(1)}(x_\ell) \right\} \\
& \quad f_N^{(1)}(dZ_1) f_N^{(1)}(dZ_2).
\end{aligned}$$

As shown in [10], the integral

$$\int \delta_1(r^N(1,2)(\mathbf{x})) \prod_{\ell=3}^N d\rho_N^{(1)}(x_\ell)$$

can be interpreted as the expectation of $\delta_1(r^N(1,2)(\mathbf{x}))$ for fixed (x_1, x_2) when x_3, \dots, x_N are drawn randomly and independently with probability $\rho_N^{(1)}$. Using the combinatorial approach of [10] to evaluate this probability, we obtain

$$\begin{aligned}
\mathbb{E}_{\rho_N^{(1)}} [\delta_1(r^N(1,2)(\mathbf{x}))] &= \sum_{R=1}^{N-1} \delta_1(R) \binom{N-2}{R-1} p^{R-1} (1-p)^{N-2-(R-1)} \\
&= (1-p)^{N-2},
\end{aligned}$$

with

$$p := M_{\rho_N^{(1)}}(x_1, |x_2 - x_1|),$$

which gives the stated result. \square

For a given smooth function ρ and $x \in \mathbb{R}^d$, let $m \in [0, 1) \mapsto R_\rho(x, m) \in [0, \infty)$ be the inverse function of $r \mapsto M_\rho(x, r)$. Note that $R_\rho(x, 0) = 0$.

Proposition 4. *For any test function $\phi(Z_1)$, we have*

$$\int \partial_t f_N^{(1)}(Z_1) \phi(Z_1) dZ_1 = \int [\phi(x_1, v_2) - \phi(Z_1)] \mathcal{F}_{f_N^{(1)}}(x_1, v_2) f_N^{(1)}(dZ_1) dv_2, \tag{10}$$

where for given smooth functions $(x, v) \mapsto \eta(x, v)$ and $\rho_\eta(x) = \int \eta(x, v) dv$, we define

$$\mathcal{F}_\eta^N(x, v) := \lambda(N) (N-1) \int_0^1 \mathcal{G}_\eta(x, v, m) (1-m)^{N-2} dm. \quad (11)$$

and

$$\mathcal{G}_\eta(x, v, m) = \frac{\int_{\mathbb{S}^{d-1}} \eta(x + R_{\rho_\eta}(x, m)\omega, v) d\omega}{\int_{\mathbb{S}^{d-1}} \rho_\eta(x + R_{\rho_\eta}(x, m)\omega) d\omega}.$$

Proof. Using polar coordinates $x_2 = x_1 + r\omega$, $r \in [0, \infty)$, $\omega \in \mathbb{S}^{d-1}$ with $\int_{\mathbb{S}^{d-1}} d\omega = 1$, we can rewrite Equation (9) as

$$\int \partial_t f_N^{(1)}(Z_1) \phi(Z_1) dZ_1 = \int [\phi(x_1, v_2) - \phi(Z_1)] \mathcal{F}_{f_N^{(1)}}^N(x_1, v_2) f_N^{(1)}(dZ_1) dv_2$$

where, we temporarily define \mathcal{F}_η^N , for a given function $(x, v) \mapsto \eta(x, v)$ by

$$\mathcal{F}_\eta^N(x, v) = \lambda(N) (N-1) \int \eta(x + r\omega, v) (1 - M_{\rho_\eta}(x, r))^{N-2} r^{d-1} dr d\omega.$$

We perform the change of variable $m := M_{\rho_\eta}(x, r)$, so that $r := R_{\rho_\eta}(x, m)$. It is then straightforward to see that \mathcal{F}_η^N is indeed defined by (11) as the jacobian of the diffeomorphism is

$$r^{d-1} dr = \frac{dm}{\int_{\mathbb{S}^{d-1}} \rho_\eta(x + R_{\rho_\eta}(x, m)\omega) d\omega}.$$

□

3.2 Limit equation

3.2.1 Preliminaries

The passage to the limit with different scaling assumptions on λ in the various considered models will use a fundamental lemma: Let a and b be two positive parameters. Define the Beta-distribution, $\beta_{a,b}$, the probability density function given for all $s \in [0, 1]$ by

$$\beta_{a,b}(s) := \frac{s^{a-1} (1-s)^{b-1}}{\mathcal{B}(a,b)},$$

where the Beta-function $\mathcal{B}(a, b)$ is defined by

$$\mathcal{B}(a, b) := \int_0^1 u^{a-1} (1-u)^{b-1} du.$$

Lemma 5 (Expected value of sequence of Beta-distribution). *Let $(h^N)_{N \in \mathbb{N}}$ be a sequence of uniformly bounded smooth functions $[0, 1] \rightarrow \mathbb{R}$ and $(b_N)_{N \in \mathbb{N}}$ be*

a sequence going to ∞ . If $h^N(u)$ converges to 0 as u goes to 0 uniformly in N then the expected value of h^N under the Beta-distribution of parameters (a, b_N) :

$$\mathbb{E}_{\beta_{a,b_N}}[h^N] = \frac{\int_0^1 h^N(u) u^{a-1} (1-u)^{b_N-1} du}{\mathcal{B}(a, b_N)}$$

converges to 0 when N goes to ∞ .

Proof. As $h^N(u)$ converges to 0 as u goes to 0 uniformly in N , we have

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } \forall u \in (0, \delta), \forall N \in \mathbb{N}, |h^N(u)| < \varepsilon.$$

Moreover, by assumption, there exists C such that for all $u \in (0, 1)$ and $N \in \mathbb{N}$, $|h^N(u)| < C$. Therefore

$$\mathbb{E}_{\beta_{a,b_N}}[h^N] \leq \varepsilon + C \frac{\int_{\delta}^1 u^{a-1} (1-u)^{b_N-1} du}{\mathcal{B}(a, b_N)} \quad (12)$$

Note that the fraction is closely related to the incomplete regularized Beta-distribution. On one hand, the function $u \mapsto u^{a-1} (1-u)^{b_N-1}$ is increasing on $[0, u_{a,b_N}]$ and decreasing on $[u_{a,b_N}, 1]$ where

$$u_{a,b_N} := \frac{a-1}{a+b_N-2},$$

is the maximum on $[0, 1]$. As $(b_N)_{N \in \mathbb{N}}$ goes to ∞ , for N large enough, u_{a,b_N} is less than δ . As a consequence, for N large enough

$$\int_{\delta}^1 u^{a-1} (1-u)^{b_N-1} du \leq \delta^{a-1} (1-\delta)^{b_N-1}. \quad (13)$$

On the other hand, by standard properties of the Beta-distribution, see [4], as N goes to ∞ and a is fixed

$$\mathcal{B}(a, b_N) = \int_0^1 u^{a-1} (1-u)^{b_N-1} du \sim \Gamma(a) b_N^{-a}, \quad (14)$$

where Γ is the Gamma-function. Collecting (13) and (14) we obtain that there exists C such that

$$\frac{\int_{\delta}^1 u^{a-1} (1-u)^{b_N-1} du}{\mathcal{B}(a, b_N)} \leq C(1-\delta)^{b_N-1} b_N^a.$$

This term converges to 0 for N large. Coming back to (12) gives the stated result. \square

3.2.2 Case $\lambda(N) = 1$

In this section, we show that the kinetic model obtained with $\lambda(N) = 1$ is trivial, *i.e.* it involves no contribution of the particle interactions to the final dynamics.

Proposition 6 (Case $\lambda(N) = 1$). *Assume that $(f_N^{(1)})_{N \in \mathbb{N}}$ and $(\rho_N^{(1)})_{N \in \mathbb{N}}$ converge toward smooth functions f and ρ respectively. If the convergence of $(f_N^{(1)})_{N \in \mathbb{N}}$ to f is such that the sequence $h^N = \mathcal{G}_{f_N^{(1)}}(x_1, v_2, \cdot) - \mathcal{G}_f(x_1, v_2, 0)$ satisfies the assumptions of Lemma 5 then we have*

$$\int \partial_t f(Z_1) \phi(Z_1) dZ_1 = 0.$$

In strong form and after restoring the drift term, the equation for f is the free transport equation:

$$\partial_t f + v \cdot \nabla_x f = 0. \quad (15)$$

Proof. By Lemma 5 applied to $\lambda = 1$, $b_N = N - 1$, we have

$$\lim_{N \rightarrow \infty} \mathcal{F}_{f_N^{(1)}}^N(x_1, v_2) = \mathcal{G}_f(x_1, v_2, 0) = \frac{f(x_1, v_2)}{\rho(x_1)}.$$

Therefore we have

$$\lim_{N \rightarrow \infty} \int \partial_t f_N^{(1)}(Z_1) \phi(Z_1) dZ_1 = \int [\phi(x_1, v_2) - \phi(x_1, v_1)] \frac{f(x_1, v_2)}{\rho(x_1)} f(dZ_1) dv_2.$$

If we exchange v_1 and v_2 , the term $\phi(x_1, v_2) - \phi(x_1, v_1)$ changes of sign while

$$f(x_1, v_1) \frac{f(x_1, v_2)}{\rho(x_1)}$$

remains unchanged, so that the integral vanishes. \square

3.2.3 Non-trivial limit

Here, we determine what must be the expression of the scaling factor $\lambda(N)$ such that the contribution of the particle interactions in the limit $N \rightarrow \infty$ is non-trivial and we determine the corresponding limit model. Before stating the main theorem, we need the following technical lemma.

Lemma 7 (Higher order expansion). *Let $(x, v) \mapsto \eta(x, v)$ be a smooth function and $\rho_\eta(x) := \int \eta(x, v) dv$. Define*

$$\mathcal{D}[\rho_\eta, \eta](x, v) = \Delta_x \eta(x, v) - \frac{\eta(x, v)}{\rho_\eta(x)} \Delta_x \rho_\eta(x). \quad (16)$$

For m small enough, we have for all $(x, v) \in \mathbb{R}^{2d}$:

$$\mathcal{G}_\eta(x, v, m) - \mathcal{G}_\eta(x, v, 0) = \frac{m^{\frac{2}{d}} d^{\frac{2}{d}-1}}{2} \frac{1}{\rho_\eta(x)^{1+\frac{2}{d}}} \mathcal{D}[\rho_\eta, \eta](x, v) + o(m^{\frac{2}{d}}), \quad (17)$$

where recall that

$$\mathcal{G}_\eta(x, v, 0) = \frac{\eta(x, v)}{\rho_\eta(x)}.$$

Proof. We need to develop $m \mapsto \mathcal{G}_\eta(x, v, m)$ for m small to higher order terms. To do so we first compute the next two orders of \mathcal{G}_η expanded in powers of $R_{\rho_\eta} = R_{\rho_\eta}(x, m)$:

$$\begin{aligned} \mathcal{G}_\eta(x, v, m) - \mathcal{G}_\eta(x, v, 0) = & \\ R_{\rho_\eta} & \left[\frac{\int (\omega \cdot \nabla_x \eta)(x + R_{\rho_\eta} \omega, v) \, d\omega}{\int \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega} \right. \\ & - \frac{\int \eta(x + R_{\rho_\eta} \omega, v) \, d\omega}{\left(\int \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \right)^2} \int \omega \cdot \nabla_x \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \left. \right] \\ & + \frac{R_{\rho_\eta}^2}{2} \left[\frac{\int (\omega \cdot \nabla_x)^2 \eta(x + R_{\rho_\eta} \omega, v) \, d\omega}{\int \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega} \right. \\ & - 2 \frac{\int (\omega \cdot \nabla_x \eta)(x + R_{\rho_\eta} \omega, v) \, d\omega}{\left(\int \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \right)^2} \int \omega \cdot \nabla_x \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \\ & + 2 \frac{\int \eta(x + R_{\rho_\eta} \omega, v) \, d\omega}{\left(\int \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \right)^3} \left(\int \omega \cdot \nabla_x \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \right)^2 \\ & \left. - \frac{\int \eta(x + R_{\rho_\eta} \omega, v) \, d\omega}{\left(\int \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \right)^2} \left(\int (\omega \cdot \nabla_x)^2 \rho_\eta(x + R_{\rho_\eta} \omega) \, d\omega \right)^2 \right] \end{aligned}$$

Fortunately all the first order terms and the second and third terms in the expression of the second order terms are zero by anti-symmetry. Now, since $\int_{\mathbb{S}^{d-1}} \omega \otimes \omega \, d\omega = \text{Id}/d$, we obtain when $R_{\rho_\eta} \rightarrow 0$:

$$\mathcal{G}_\eta(x, v, m) - \mathcal{G}_\eta(x, v, 0) = \frac{R_{\rho_\eta}^2}{2d} \left[\frac{\Delta \eta(x, v)}{\rho_\eta(x)} - \frac{\eta(x, v) \Delta_x \rho_\eta(x)}{\rho_\eta(x)^2} \right] + o\left(R_{\rho_\eta}^2\right). \quad (18)$$

Let us now develop $m \mapsto R_{\rho_\eta}(x, m)$ in powers of m . We have for r small enough

$$M(x, r) := \int_{|x-x'| < r} \rho_\eta(x') \, dx' = \rho_\eta(x) \frac{r^d}{d} + o(r^d).$$

Therefore for m small enough

$$R_{\rho_\eta} = \left(\frac{dm}{\rho_\eta(x)} \right)^{\frac{1}{d}} + o(m^{\frac{1}{d}}),$$

which concludes the proof by inserting this expression in (18). \square

Before stating the theorem, we introduce the following notations: define

$$h^N(m) = H^N(m) - H(0), \quad (19)$$

where

$$H^N(m) := \frac{\mathcal{G}_{f_N^{(1)}}(x_1, v_2, m) - \mathcal{G}_{f_N^{(1)}}(x_1, v_2, 0)}{m^{\frac{2}{d}}}. \quad (20)$$

and

$$H(0) := \frac{d^{\frac{2}{d}-1}}{2} \frac{1}{\rho(x_1)^{\frac{2}{d}+1}} \mathcal{D}[\rho, f](x_1, v_2). \quad (21)$$

We note that from Lemma 7, $h^N(m) \rightarrow 0$ as $m \rightarrow 0$, for all integers N . Now we state the main theorem of this paper:

Theorem 8. *Assume that $(f_N^{(1)})_{N \in \mathbb{N}}$ and $(\rho_N^{(1)})_{N \in \mathbb{N}}$ converge toward smooth functions f and ρ respectively. Assume that the convergence of $(f_N^{(1)})_{N \in \mathbb{N}}$ to f is such that the sequence h^N defined by (19) satisfies the assumptions of Lemma 5. Set $\lambda(N)$ to the value*

$$\lambda(N) = \frac{1}{(N-1) \frac{d^{\frac{2}{d}-1}}{2} \mathcal{B}(1 + \frac{2}{d}, N-1)}. \quad (22)$$

Then for all test functions $\phi(Z_1)$ we have

$$\begin{aligned} & \int \partial_t f(Z_1) \phi(Z_1) \, dZ_1 \\ &= \int \phi(x_1, v_2) \frac{1}{\rho(x_1)^{\frac{2}{d}+1}} \left[\Delta_x f(Z_1) - \frac{f(Z_1)}{\rho(x_1)} \Delta_x \rho(x_1) \right] \, dZ_1 \, dv_2. \end{aligned} \quad (23)$$

In strong form, introducing the collision operator

$$Q[f] := \frac{1}{\rho^{\frac{2}{d}+1}} \left[\Delta_x f - \frac{f}{\rho} \Delta_x \rho \right], \quad (24)$$

and after restoring the drift term, the equation for f reads:

$$\partial_t f + v \cdot \nabla_x f = Q[f]. \quad (25)$$

Before proving this theorem, we first prove the following intermediate step:

Proposition 9. *Under the assumptions of Theorem 8, for all test functions $\phi(Z_1)$, we have*

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_0^1 \left[\mathcal{G}_{f_N^{(1)}}(x_1, v_2, m) - \mathcal{G}_{f_N^{(1)}}(x_1, v_2, 0) \right] \frac{(1-m)^{N-2}}{\mathcal{B}(1 + \frac{2}{d}, N-1)} \, dm \\ = \frac{d^{\frac{2}{d}-1}}{2} \frac{1}{\rho(x_1)^{\frac{2}{d}+1}} \mathcal{D}[\rho, f](x_1, v_2). \end{aligned} \quad (26)$$

Proof of Proposition 9. Thanks to the assumptions on h^N , we can apply Lemma 5 with $a = 1 + \frac{2}{d}$ and $b_N = N - 1$, and obtain

$$\lim_{N \rightarrow \infty} \int_0^1 h^N(m) d\beta_{\frac{2}{d}-1, N-1} = 0.$$

As $\beta_{\frac{2}{d}-1, N-1}$ is a probability this proves that

$$\lim_{N \rightarrow \infty} \int_0^1 H^N(m) d\beta_{1+\frac{2}{d}, N-1} = H(0). \quad (27)$$

Using (20), we have

$$\begin{aligned} & \int_0^1 H^N(m) d\beta_{1+\frac{2}{d}, N-1} \\ &= \int_0^1 \left[\mathcal{G}_{f_N^{(1)}}(x_1, v_2, m) - \mathcal{G}_{f_N^{(1)}}(x_1, v_2, 0) \right] \frac{(1-m)^{N-2}}{\mathcal{B}(1+\frac{2}{d}, N-1)} dm. \end{aligned} \quad (28)$$

Inserting (28) and the expression of $H(0)$ given by (21) into (27) leads to (26). \square

Proof of Theorem 8. We start from (10) where \mathcal{F}^N is defined by (11). Using (26) and owing to the fact that the term involving $\mathcal{G}_{f_N^{(1)}}(x_1, v_2, 0)$ vanishes upon integration with respect to (Z_1, v_2) (exactly like in the proof of Proposition 6), we deduce that, for N large, up to negligible terms when $N \rightarrow \infty$, we have

$$\begin{aligned} & \int \partial_t f(Z_1) \phi(Z_1) dZ_1 \\ &= \lambda(N) (N-1) \frac{d^{\frac{2}{d}-1}}{2} \mathcal{B}(1+\frac{2}{d}, N-1) \\ & \int [\phi(x_1, v_2) - \phi(x_1, v_1)] \frac{f(Z_1)}{\rho(x_1)^{\frac{2}{d}+1}} \mathcal{D}[\rho, f](x_1, v_2) dZ_1 dv_2. \end{aligned}$$

Using Definition (16) of \mathcal{D} , we see that the second term of $f(Z_1) \mathcal{D}[\rho, f](x_1, v_2)$ is symmetric by exchange of (v_1, v_2) . Since $[\phi(x_1, v_2) - \phi(x_1, v_1)]$ is anti-symmetric under this exchange, this second term vanishes after integration. Therefore, the only non-zero term comes from the first term of \mathcal{D} , which leads to (up to negligible terms when $N \rightarrow \infty$) to

$$\begin{aligned} & \int \partial_t f(Z_1) \phi(Z_1) dZ_1 \\ &= \lambda(N) (N-1) \frac{d^{\frac{2}{d}-1}}{2} \mathcal{B}(1+\frac{2}{d}, N-1) \\ & \int \phi(x_1, v_2) \frac{1}{\rho(x_1)^{\frac{2}{d}+1}} \left[\Delta_x f(Z_1) - \frac{f(Z_1)}{\rho(x_1)} \Delta_x \rho(x_1) \right] dZ_1 dv_2, \end{aligned}$$

and with the choice (22), this leads to (23). It readily seen, applying Green's formula, that the strong form of the equation is (25), upon inserting back the drift term. \square

Remark 10. We note that $\lambda(N) \rightarrow \infty$ as $N \rightarrow \infty$. In particular, in the case $d = 2$, $\lambda(N)$ can be easily computed and has value $\lambda(N) = 2N$. This means that the limit is non-trivial only if the number of collisions per unit time is larger than for the standard kinetic scale (which corresponds to $\lambda(N) = 1$).

4 K -nearest neighbor dynamics

4.1 Master equation

Let K be a fixed integer in $\{1, \dots, N\}$ and a sequence $(\alpha_k)_{k \in \{1, \dots, K\}}$ such that $\sum_{k=1}^K \alpha_k = 1$ being given. We extend this sequence for k in $\{1, \dots, N\}$ by taking $\alpha_k = 0$ for all $k > K$. The particle i will adopt the velocity of its j th-nearest neighbor with a probability π_{ij}^N given by (7). Section 3 was devoted to the case $\alpha_1 = 1$ and $\alpha_j \equiv 0$ for all $j > 1$.

As shown in [10], when the collisions occur at Poisson times with rate $\lambda(N)N$, the master equation of the N -particle distribution function $f^{(N)}$ in weak form is, for all test function $\phi^N : \mathbf{Z} \mapsto \phi^N(\mathbf{Z})$:

$$\begin{aligned} \int \partial_t f^{(N)}(\mathbf{Z}) \phi^N(\mathbf{Z}) d\mathbf{Z} &= \sum_{i=1}^N \int f^{(N)}(\mathbf{Z}) (v_i \cdot \nabla_{x_i}) \phi^N(\mathbf{Z}) d\mathbf{Z} \\ &+ \lambda(N) \sum_{i=1}^N \sum_{\substack{j=1 \\ j \neq i}}^N \iint \sum_{k=1}^K \alpha_k \delta_k[r^N(i, j)(\mathbf{x})] [\phi^N(Z_1, \dots, x_i, v_j, \dots, x_j, v_j, \dots, Z_N) \\ &\quad - \phi^N(\mathbf{Z})] f^{(N)}(\mathbf{Z}) d\mathbf{Z}. \end{aligned} \quad (29)$$

4.2 Propagation of chaos

Proposition 11 (First marginal equation with propagation of chaos). *Under the propagation of chaos assumption (2) we have for all test function ϕ*

$$\begin{aligned} \frac{1}{\lambda(N)(N-1)} \int \partial_t f_N^{(1)}(Z_1) \phi(Z_1) dZ_1 &= \sum_{k=1}^K \alpha_k \binom{N-2}{k-1} \\ \int [\phi(x_1, v_2) - \phi(Z_1)] M_{\rho_N^{(1)}}(x_1, |x_2 - x_1|)^{k-1} &\left(1 - M_{\rho_N^{(1)}}(x_1, |x_2 - x_1|)\right)^{N-k-1} \\ &f_N^{(1)}(dZ_1) f_N^{(1)}(dZ_2). \end{aligned} \quad (30)$$

Proof. As before we have

$$\begin{aligned} & \frac{1}{\lambda(N)(N-1)} \int \partial_t f^{(N)}(\mathbf{Z}) \phi^N(Z_1) \, d\mathbf{Z} \\ &= \sum_{k=1}^K \alpha_k \int \sum_{k=1}^K \delta_k[r^N(1,2)(\mathbf{x})] [\phi(x_1, v_2) - \phi(Z_1)] f^{(N)}(\mathbf{Z}) \, d\mathbf{Z}, \end{aligned}$$

which leads, by the propagation of chaos assumption (2) to

$$\begin{aligned} & \frac{1}{\lambda(N)(N-1)} \int \partial_t f_N^{(1)}(\mathbf{Z}) \phi(Z_1) \, d\mathbf{Z} \\ &= \sum_{k=1}^K \alpha_k \int [\phi(x_1, v_2) - \phi(Z_1)] \left\{ \int \delta_k[r^N(1,2)(\mathbf{x})] \prod_{\ell=3}^N d\rho_N^{(1)}(x_\ell) \right\} \\ & \qquad \qquad \qquad f_N^{(1)}(dZ_1) f_N^{(1)}(dZ_2), \end{aligned}$$

up to negligible terms as $N \rightarrow \infty$. The computation of the integral

$$\int \delta_k(r^N(1,2)(\mathbf{x})) \prod_{\ell=3}^N d\rho_N^{(1)}(x_\ell)$$

is slightly different from before as here (see [10] for details)

$$\begin{aligned} \mathbb{E}_{\rho_N^{(1)}} [\delta_k(r^N(1,2)(\mathbf{x}))] &= \sum_{R=1}^{N-1} \delta_k(R) \binom{N-2}{R-1} p^{R-1} (1-p)^{N-2-(R-1)} \\ &= \binom{N-2}{k-1} p^{k-1} (1-p)^{N-k-1}, \end{aligned}$$

with

$$p := M_{\rho_N^{(1)}}(x_1, |x_2 - x_1|),$$

which gives the stated result. \square

The probability considered in Section 3 was the Bernoulli distribution. A similar result can be proved with a more general binomial distribution: For a given smooth function ρ and $x \in \mathbb{R}^d$, let $m \in [0, 1] \mapsto R_\rho(x, m) \in [0, \infty)$ be the inverse function of $r \mapsto M_\rho(x, r)$. We also define the binomial distribution with parameters $n \in \mathbb{N}$ and $p \in [0, 1]$, denoted $\mathcal{M}(n, p)$, as the probability on the discrete set $\{0, 1, \dots, N\}$ given by the probability mass function:

$$\mu(k; n, p) = \mathbb{P}(X = k) = \binom{n}{k} p^k (1-p)^{n-k}.$$

We note that, by integration by parts, we have

$$(N-1) \int_0^1 \mu(k; n, p) \, dp = 1. \quad (31)$$

Proposition 12. *We have*

$$\begin{aligned} \int \partial_t f_N^{(1)}(Z_1) \phi(Z_1) dZ_1 \\ = \int [\phi(x_1, v_2) - \phi(Z_1)] \mathcal{F}_{f_N^{(1)}, K}^N(x_1, v_2) f_N^{(1)}(dZ_1) dv_2 \end{aligned}$$

where for given smooth functions $(x, v) \mapsto \eta(x, v)$ and $\rho_\eta(x) = \int \eta(x, v) dv$, we define

$$\mathcal{F}_{\eta, K}^N(x, v) := \lambda(N) \int_0^1 \mathcal{G}_\eta(x, v, m) \nu_K(m) dm. \quad (32)$$

with

$$\nu_K(m) := (N-1) \sum_{k=1}^K \alpha_k \mu(k-1; N-2, m)$$

and

$$\mathcal{G}_\eta(x, v, m) = \frac{\int_{\mathbb{S}^{d-1}} \eta(x + R_{\rho_\eta}(x, m)\omega, v) d\omega}{\int_{\mathbb{S}^{d-1}} \rho_\eta(x + R_{\rho_\eta}(x, m)\omega) d\omega}.$$

The proof is identical and will not be repeated here. Also note that $\nu_K(m)$, dm is a probability on $[0, 1]$. Indeed, thanks to (31), we have

$$\int_0^1 \nu_K(m) dm = (N-1) \sum_{k=1}^K \alpha_k \int \mu(k-1; N-2, m) dm = \sum_{k=1}^K \alpha_k = 1.$$

4.3 Limit equation

4.3.1 Case $\lambda(N) = 1$

Proposition 13 (Case $\lambda(N) = 1$). *Assume that $(f_N^{(1)})_{N \in \mathbb{N}}$ and $(\rho_N^{(1)})_{N \in \mathbb{N}}$ converge toward smooth functions f and ρ respectively. If the convergence of $(f_N^{(1)})_{n \in \mathbb{N}}$ to f is such that the sequence $h^N = \mathcal{G}_{f_N^{(1)}}(x_1, v_2, \cdot) - \mathcal{G}_f(x_1, v_2, 0)$ satisfies the assumptions of Lemma 5 then for all test functions ϕ we have*

$$\int \partial_t f(Z_1) \phi(Z_1) dZ_1 = 0.$$

In strong form and after restoring the drift term, the equation for f is the free transport equation (15).

Proof. As $\nu_K(m)$ is a convex linear combination of binomial distributions, we can apply Lemma 5 to $\mu(k-1; N-2, m)$ for any k and this leads to the result. \square

4.3.2 Non-trivial case

Like in section 3.2.3, we now determine the scaling factor $\lambda(N)$ for the particle interactions to have a non-trivial contribution in the limit $N \rightarrow \infty$ and we determine the corresponding limit model.

Theorem 14. Assume that $(f_N^{(1)})_{N \in \mathbb{N}}$ and $(\rho_N^{(1)})_{N \in \mathbb{N}}$ converge toward smooth functions f and ρ respectively. Assume that the convergence of $(f_N^{(1)})_{N \in \mathbb{N}}$ to f is such that the sequence h^N defined by (19) satisfies the assumptions of Lemma 5. Set $\lambda(N)$ to the value

$$\lambda(N) = \frac{1}{(N-1) \frac{d^{\frac{2}{d}-1}}{2} \sum_{k=1}^K \alpha_k \binom{N-2}{k-1} \mathcal{B}(k + \frac{2}{d}, N-k)}. \quad (33)$$

Then for all test functions $\phi(Z_1)$, f satisfies Eq. (23). In strong form and after restoring the drift term, the equation for f is given by (25)

The result still relies on Lemma 7 and the following analog of Proposition 9:

Proposition 15. Under the assumptions of Theorem 14, for all test functions $\phi(Z_1)$, we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_0^1 \left[\mathcal{G}_{f_N^{(1)}}(x_1, v_2, m) - \mathcal{G}_{f_N^{(1)}}(x_1, v_2, 0) \right] \\ \times \frac{\sum_{k=1}^K \alpha_k \binom{N-2}{k-1} m^{k-1} (1-m)^{N-k-1}}{\sum_{k=1}^K \alpha_k \binom{N-2}{k-1} \mathcal{B}(k + \frac{2}{d}, N-k)} dm \\ = \frac{d^{\frac{2}{d}-1}}{2} \frac{1}{\rho(x_1)^{\frac{2}{d}+1}} \mathcal{D}[\rho, f](x_1, v_2) \end{aligned}$$

The proof Proposition 15 is similar to that of Proposition 9 and is left to the reader. The deduction of Theorem 14 from Proposition 15 is exactly the same as that of Theorem 8 from Proposition 9.

Remark 16. We note the following property which can be proven by integration by parts: for all $k \in \mathbb{Z}$, for all $a \in [-k, \infty)$, for all $b \in [k, \infty)$, we have:

$$\mathcal{B}(a+k, b-k) = \mathcal{B}(a, b) \binom{a+k-1}{k} \binom{b-1}{k}^{-1}.$$

Thanks to this property, we have for $k \in \mathbb{N}$, $k \geq 2$:

$$\binom{N-2}{k-1} \mathcal{B}(k + \frac{2}{d}, N-k) = \binom{\frac{2}{d} + k - 1}{k-1} \mathcal{B}(1 + \frac{2}{d}, N-1).$$

Thus,

$$\lambda(N) = \frac{1}{\sum_{k=1}^K \binom{\frac{2}{d} + k - 1}{k-1} \alpha_k} \frac{1}{(N-1) \frac{d^{\frac{2}{d}-1}}{2} \mathcal{B}(1 + \frac{2}{d}, N-1)}.$$

Therefore, we find the same scaling of $\lambda(N)$ as in the nearest neighbor interaction case, up to the multiplication by the constant factor $(\sum_{k=1}^K \binom{\frac{2}{d} + k - 1}{k-1} \alpha_k)^{-1}$.

5 From smooth to nearest-neighbor interaction

In this section we investigate the connection between the smooth rank-based interaction developed in [10] and the nearest neighbor (or K nearest-neighbor) interaction considered here. We show that when the kernel $K(m)$ concentrates near $m = 0$, we pass from (3) to (5). As the kernel K concentrates, it needs to be rescaled in the appropriate way. More precisely, we introduce a rescaling parameter ε which we will tend to zero and the following rescaled kernel

$$K^\varepsilon(m) = \frac{1}{\varepsilon^{1+\frac{2}{d}}} K^0\left(\frac{m}{\varepsilon}\right), \quad (34)$$

and we assume that K_0 is normalized such that

$$\frac{d^{\frac{2}{d}-1}}{2} \int m^{\frac{2}{d}} K^0(m) dm = 1. \quad (35)$$

Proposition 17. *The kinetic nearest-neighbor interaction model (5) is the limit when ε goes to 0 of the kinetic smooth rank-based interaction model (3) with interaction kernel $K = K^\varepsilon$ given by (34) with normalization given by (35).*

Proof. The weak form of the kinetic smooth rank-based interaction model (3) obtained in [10] is given, for any test function $\phi(Z_1)$ by

$$\int \partial_t f(Z_1, t) \phi(Z_1) dZ_1 = \int [\phi(Z_1) - \phi(x_1, v_2)] K^\varepsilon(M_\rho(x_1, |x_2 - x_1|)) f(dZ_1) f(dZ_2).$$

Passing to polar coordinates in x_2 , i.e. writing $x_2 = x_1 + r\omega$, $r = |x_2 - x_1|$, $\omega = \frac{x_2 - x_1}{|x_2 - x_1|}$, this equation reads

$$\int \partial_t f(Z_1, t) \phi(Z_1) dZ_1 = \int [\phi(Z_1) - \phi(x_1, v_2)] f(x_1 + r\omega, v_2) K^\varepsilon(M_\rho(x_1, r)) f(dZ_1) r^{d-1} dr d\omega.$$

Using the change of coordinates $m = M_\rho(x_1, r)$ the inverse function of which is $r = R_\rho(x_1, m)$ and the jacobian of which is

$$dm = \rho(x_1 + r\omega) r^{d-1} dr,$$

we obtain

$$\int \partial_t f(Z_1, t) \phi(Z_1) dZ_1 = \int [\phi(Z_1) - \phi(x_1, v_2)] \mathcal{G}(x_1, v_2, m) K^\varepsilon(m) f(dZ_1) dm d\omega dv_2.$$

where

$$\mathcal{G}(x, v, m) := \frac{f(x + R_\rho(x, m)\omega, v)}{\rho(x_1 + R_\rho(x_1, m)\omega)}.$$

Let us look at the limit of $\int K^\varepsilon(m)\mathcal{G}(x, v, m) dm$ when ε goes to 0. By Lemma 7, we have for small ε

$$\begin{aligned} \int K^\varepsilon(m)\mathcal{G}(x, v, m) dm &= \frac{1}{\varepsilon^{1+\frac{2}{d}}} \int K^0\left(\frac{m}{\varepsilon}\right) \mathcal{G}(m) dm = \frac{1}{\varepsilon^{\frac{2}{d}}} \int K^0(m) \mathcal{G}(\varepsilon m) dm \\ &= \frac{1}{\varepsilon^{\frac{2}{d}}} \int K^0(m) \left[\mathcal{G}(x, v, 0) + \frac{\varepsilon^{\frac{2}{d}}}{2} m^{\frac{2}{d}} \mathcal{G}^{(2)}(x, v, 0) + o(\varepsilon^{\frac{2}{d}}) \right] dm \end{aligned}$$

where by (17)

$$\mathcal{G}^{(2)}(x, v, 0) := \frac{d^{\frac{2}{d}-1}}{2} \frac{1}{\rho(x)^{\frac{2}{d}+1}} \mathcal{D}[\rho, f](x, v),$$

As the first term is anti-symmetric in the transform $(v_1, v_2) \rightarrow (v_2, v_1)$, we obtain at the limit $\varepsilon \rightarrow 0$ and thanks again to Lemma 7:

$$\begin{aligned} &\int \partial_t f(Z_1, t) \phi(Z_1) dZ_1 \\ &= \int m^{\frac{2}{d}} K^0(m) dm \int [\phi(Z_1) - \phi(x_1, v_2)] \mathcal{G}^{(2)}(x_1, v_2, 0) f(dZ_1) dv_2 \\ &= \left(\frac{d^{\frac{2}{d}-1}}{2} \int m^{\frac{2}{d}} K^0(m) dm \right) \int [\phi(Z_1) - \phi(x_1, v_2)] \frac{1}{\rho(x_1)^{\frac{2}{d}+1}} \\ &\quad \mathcal{D}[\rho, f](x_1, v_2) f(dZ_1) dv_2. \end{aligned}$$

The passage to the strong form is identical to that of Section 3.2.3 and we obtain

$$\partial_t f = \left(\frac{d^{\frac{2}{d}-1}}{2} \int m^{\frac{2}{d}} K^0(m) dm \right) Q[f],$$

where Q is defined by (24). This concludes the proof assuming that K_0 is normalized according to (35). \square

6 Conclusion

In this paper, we have put forward a particle interaction model where particles interact with their nearest neighbor. We have shown that the large particle limit under the Propagation of Chaos assumption is a spatial diffusion for the particle distribution function corrected by an anti-diffusion term acting on the spatial density. We have shown that the appearance of this anti-diffusion term depending on the spatial density results from the fact that the interactions are mass-preserving. We have also considered a model in which particles interact with their K nearest neighbors, for a fixed value of K , showing that the corresponding kinetic model is the same as in the nearest neighbor interaction case. Finally, we have linked this work with the previous article [10]

where smooth rank-based dynamics were considered and shown that the kinetic nearest-neighbor model can be recovered from the former through a singular limit involving a scaling of the interaction kernel.

The kinetic models obtained here, as in [10], are novel. Their mathematical theory is entirely open: proving existence and uniqueness of solutions, investigating large-time behavior, equilibria and other qualitative properties of the solutions will require the establishment of an appropriate mathematical framework. In parallel, more elaborate physical interaction models should also be considered. One important question is to investigate how the present results are robust to the introduction of noise in the interaction dynamics. Indeed, noise play an important part of many flocking models in nature. Finally, adequate numerical methods for the kinetic models must be developed and the assessment of the kinetic models against the particle ones in realistic situations should be carefully documented so that these models can be used in practice.

Acknowledgements: PD acknowledges support by the Engineering and Physical Sciences Research Council (EPSRC) under grants no. EP/M006883/1, EP/N014529/1 and EP/P013651/1, by the Royal Society and the Wolfson Foundation through a Royal Society Wolfson Research Merit Award no. WM130048 and by the National Science Foundation (NSF) under grant no. RNMS11-07444 (KI-Net). PD is on leave from CNRS, Institut de Mathématiques de Toulouse, France. AB acknowledges support from the IDEX-UNITI-ANR-11-IDEX-0002-Transversalité-Projet MUSE "Multi-disciplinary study of emergence phenomena".

Data statement: no new data were collected in the course of this research.

References

- [1] Albi, G., Balagué, D., Carrillo, J. A., Von Brecht, J.: Stability analysis of flock and mill rings for second order models in swarming. *SIAM J. Appl. Math.* 74, 794-818 (2014)
- [2] Albi, G., Pareschi, L.: Binary interaction algorithms for the simulation of flocking and swarming dynamics. *Multiscale Model. Simul.* 11, 1-29 (2013)
- [3] Aoki, I.: A simulation study on the schooling mechanism in fish. *Bulletin of the Japan Society of Scientific Fisheries* 48, 1081–1088 (1982)
- [4] Askey, R., Roy, R.: Gamma function, *NIST handbook of mathematical functions*. US Dept. Commerce, Washington DC, 135–147 (2010)
- [5] Ballerini, M., Cabibbo, N., Candelier, R., Cavagna, A., Cisbani, E., Giardina, I., Lecomte, V., Orlandi, A., Parisi, G., Procaccini, A., Viale, M., Zdravkovic, V.: Interaction ruling animal collective behavior depends on topological rather than metric distance: Evidence from a field study. *Proc. Natl. Acad. Sci. USA* 105, 1232-1237 (2008)

- [6] Banner, A. D., Fernholz, R., Karatzas, I.: Atlas models of equity markets. *Ann. Appl. Probab.* 15, 2296-2330 (2005)
- [7] Barbaro, A. B., Cañizo, J. A., Carrillo, J. A., Degond, P.: Phase Transitions in a kinetic flocking model of Cucker-Smale type. *Multiscale Model. Simul.* 14, 1063-1088 (2016)
- [8] Bellomo, N., Soler, J.: On the mathematical theory of the dynamics of swarms viewed as complex systems. *Math. Models Methods Appl. Sci.* 22 supp01, 1140006 (2012)
- [9] Bertozzi, A. L., Kolokolnikov, T., Sun, H., Uminsky, D., Von Brecht, J.: Ring patterns and their bifurcations in a nonlocal model of biological swarms. *Commun. Math. Sci.* 13, 955-985 (2015)
- [10] Blanchet, A., Degond, P.: Topological interactions in a Boltzmann-type framework. *J. Stat. Phys.* 163, 41-60 (2016)
- [11] Bode, N. W., Franks, D. W., Wood, A. J.: Limited interactions in flocks: relating model simulations to empirical data. *Journal of The Royal Society Interface* rsif20100397 (2010)
- [12] Bolley, F., Cañizo, J. A., Carrillo, J.: A Mean-field limit for the stochastic Vicsek model. *Appl. Math. Lett.* 25, 339-343 (2012)
- [13] Brenier, Y.: Rearrangement, convection, convexity and entropy. *Philos. Trans. A* 371, 20120343 (2013)
- [14] Camperi, M., Cavagna, A., Giardina, I., Parisi, G., Silvestri, E.: Spatially balanced topological interaction grants optimal cohesion in flocking models. *Interface focus* 2, 715-725 (2012)
- [15] Carlen, E., Chatelin, R., Degond, P., Wennberg, B.: Kinetic hierarchy, propagation of chaos in biological swarm models. *Phys. D* 260, 90111 (2013)
- [16] Carlen, E., Degond, P., Wennberg, B.: Kinetic limits for pair-interaction driven master equations, biological swarm models. *Math. Models Methods Appl. Sci.* 23, 13391376 (2013)
- [17] Carrillo, J. A., Fornasier, M., Rosado, J., Toscani, G.: Asymptotic flocking dynamics for the kinetic Cucker-Smale model. *SIAM J. Math. Anal.* 42, 218-236 (2010)
- [18] Cavagna, A., Cimarelli, A., Giardina, I., Parisi, G., Santagati, R., Stefanini, F., Tavarone, R.: From empirical data to inter-individual interactions: unveiling the rules of collective animal behavior. *Math. Models Methods Appl. Sci.* 20 Suppl. 01, 1491-1510 (2010)
- [19] Degond, P., Frouvelle, A., Liu, J. G.: Macroscopic limits and phase transition in a system of self-propelled particles. *J. Nonlinear Sci.* 23, 427-456 (2013)

- [20] Degond, P., Frouvelle, A., Liu, J. G.: Phase transitions, hysteresis, and hyperbolicity for self-organized alignment dynamics. *Arch. Ration. Mech. Anal.* 216, 63-115 (2015)
- [21] Degond, P., Motsch, S.: Continuum limit of self-driven particles with orientation interaction. *Math. Models Methods Appl. Sci.* 18 supp01, 1193-1215 (2008)
- [22] Davis, D., Holt, C.: *Experimental economics*, Princeton university press, 1993
- [23] Eftimie, R.: Hyperbolic and kinetic models for self-organized biological aggregations and movement: a brief review. *J. Math. Biol.* 65, 35-75 (2012)
- [24] Eriksson, A., Jacobi, M. N., Nyström, J., Tunström, K.: Determining interaction rules in animal swarms. *Behavioral Ecology* 21, 1106-1111 (2010)
- [25] Fernholz, E. R.: *Stochastic Portfolio Theory*. Springer, New York, (2002).
- [26] Fornasier, M., Haskovec, J., Toscani, G.: Fluid dynamic description of flocking via the PovznerBoltzmann equation. *Phys. D* 240, 21-31 (2011)
- [27] Gamba, I. M., Haack, J. R., Motsch, S.: Spectral method for a kinetic swarming model. *J. Comput. Phys.* 297, 32-46 (2015)
- [28] Ginelli, F., Chaté, H.: Relevance of metric-free interactions in flocking phenomena. *Phys. Rev. Lett.* 105, 168103 (2010)
- [29] Ha, S. Y., Liu, J. G.: A simple proof of the Cucker-Smale flocking dynamics and mean-field limit. *Comm. Math. Sci.* 7, 297-325 (2009)
- [30] Ha, S. Y., Tadmor, E.: From particle to kinetic and hydrodynamic descriptions of flocking. *Kinet. Relat. Models* 1, 415-435 (2008)
- [31] Haskovec, J.: Flocking dynamics and mean-field limit in the cuckersmale-type model with topological interactions. *Phys D* 261, 42-51 (2013)
- [32] Helbing, D., Molnar, P.: Social force model for pedestrian dynamics. *Phys. Rev. E* 51, 4282-4286 (1995)
- [33] Hemelrijk, C. K., Hildenbrandt, H.: Some causes of the variable shape of flocks of birds. *PloS one* 6, e22479 (2011)
- [34] Hill, S., Buss, D.: Risk and relative social rank: positional concerns and risky shifts in probabilistic decision-making. *Evolution and Human Behavior* 31, 219-226 (2010)
- [35] Ichiba, T., Papathanakos, V., Banner, A., Karatzas, I., Fernholz, R.: Hybrid atlas models. *Ann. Appl. Probab.* 21, 609-644 (2011)

- [36] Jian, M., Wei-Guo, S., Guang-Xuan, L.: Multi-grid simulation of pedestrian counter flow with topological interaction. *Chinese physics B* 19, 128901 (2010)
- [37] Karper, T. K., Mellet, A., Trivisa, K.: Hydrodynamic limit of the kinetic CuckerSmale flocking model. *Math. Models Methods Appl. Sci.* 25, 131-163 (2015)
- [38] Lukeman, R., Li, Y. X., Edelstein-Keshet, L.: Inferring individual rules from collective behavior. *Proc. Natl. Acad. Sci. USA* 107, 12576-12580 (2010)
- [39] Martin, S.: Multi-agent flocking under topological interactions. *Systems Control Lett.* 69, 53-61 (2014)
- [40] Motsch, S., Tadmor, E.: A new model for self-organized dynamics and its flocking behavior. *J. Stat. Phys.* 144, 923-947 (2011)
- [41] Niizato, T., Murakami, H., Gunji, Y. P.: Emergence of the scale-invariant proportion in a flock from the metric-topological interaction. *Biosystems* 119, 62-68 (2014)
- [42] Norman, D.: *Memory and attention*, John Wiley and Sons New York, 1976
- [43] Parisot, M., Lachowicz, M.: A Kinetic Model for the formation of Swarms with nonlinear interactions. *Kinet. Relat. Models* 9, 131-164 (2016)
- [44] Shang, Y., Bouffanais, R.: Consensus reaching in swarms ruled by a hybrid metric-topological distance. *Eur. Phys. J. B* 87, 1-7 (2014)
- [45] Shang, Y., Bouffanais, R.: Influence of the number of topologically interacting neighbors on swarm dynamics. *Scientific reports* 4, 4184 (2014)
- [46] Wang, L., Chen, G.: Synchronization of multi-agent systems with metric-topological interactions. *Chaos* 26, 094809 (2016)